

Risk Factors Of ACS In Young Adults: A Cross-Sectional Analysis

Md. Akhwanur Rahman Chowdhury^{1*}, Md. Mojib Uddin²,
Md. Raihan Monzoor³ Md. Harun-Or-Rashid⁴, Nasrin Jahan Shammi⁵,
Md. Abul Kalam Azad⁶

Seniour Consultant, 250 Baded General Hospital, Naogaon, Bangladesh

Associate Professor, Department Of Pharmacology & Therapeutic, Rajshahi Medical College, Rajshahi, Bangladesh.

Assistant Professor, Department Of Community Medicine, Barind Medical College, Rajshahi, Bangladesh. Dr. Md. Harun-Or-Rashid.

Assistant Professor, Department Of Otorhinolaryngology, Pabna Medical College, Pabna, Bangladesh.

Professor, Department Of Pharmacology & Therapeutic, Barind Medical College, Rajshahi, Bangladesh.

Assistant Professor, Department Of Medicine, Rangpur Medical College, Rangpur, Bangladesh

Abstract

Introduction: Acute coronary syndrome (ACS) is a leading cause of morbidity and mortality worldwide, with its prevalence rising among younger adults. While traditionally considered a condition affecting older individuals, recent studies have highlighted an increasing incidence of ACS in younger populations, emphasizing the importance of identifying early risk factors. This study aimed to assess the prevalence of risk factors of ACS in young adults.

Methods: This cross-sectional observational study was conducted at the Department of Cardiology in 250 Baded General Hospital, Naogaon, Bangladesh. The study duration was one year. The study population consisted of a total of 100 young adults under the age of 35 years who were admitted to the study hospital for some form of acute coronary syndromes and had undergone coronary angiograms during the study period. Data analysis was carried out by using the statistical package SPSS version 23.0 Windows software.

Result: The study of 100 participants found that 90% were male, with the majority (67%) aged 31-35 years. Most had normal BMI (83%), and 63% reported no stress. Chest pain was the most common symptom (95%), and hypertension (70%) was the most prevalent risk factor, followed by diabetes (55%) and smoking (43%). ECG findings showed that 50% had ST-segment elevation. Lipid profiles revealed elevated triglycerides (212.0 mg/dl) and LDL-C (138.3 mg/dl), with HDL-C within normal limits (39.1 mg/dl).

Conclusion: This study concludes that the increasing prevalence of ACS among young adults and emphasizes the role of traditional risk factors such as hypertension, diabetes, dyslipidemia, smoking, and stress. The clinical and diagnostic findings in this study, including typical chest pain, ST-segment elevation on ECG, and elevated lipid levels, align with previous research, underlining the need for early detection and intervention to reduce the burden of ACS in younger populations.

Keywords: Acute Coronary Syndrome (ACS), Young adults, Risk factors, Hypertension

Date of Submission: 10-12-2024

Date of Acceptance: 20-12-2024

I. Introduction

Acute Coronary Syndrome (ACS), which includes conditions like unstable angina and myocardial infarction (MI), is a leading cause of morbidity and mortality worldwide. While ACS has traditionally been viewed as a disease affecting older adults, an increasing number of young adults are being diagnosed with cardiovascular diseases, including ACS. Recent studies indicate that ACS is becoming more prevalent in younger populations, which challenges the long-standing assumption that coronary artery disease (CAD) is primarily a concern for older individuals (1). In young adults, poor dietary habits, lack of physical activity, and rising obesity rates contribute to an increase in lipid abnormalities, placing individuals at higher risk of early-onset cardiovascular disease. Research has shown that young adults with dyslipidemia are at a significantly higher risk of developing ACS compared to those with normal lipid profiles (2). Smoking is a well-known and modifiable risk factor for ACS. The harmful chemicals in tobacco smoke cause endothelial dysfunction, increase clotting, and accelerate atherosclerosis (3). Additionally, smoking reduces the levels of HDL-C, thus exacerbating the lipid abnormalities that contribute to ACS (4). Smoking is especially prevalent among younger populations, and its

impact on cardiovascular health is seen at an earlier age compared to other risk factors. Researchers found that young adults who smoked were significantly more likely to experience a myocardial infarction than non-smokers(4). Diabetes mellitus, particularly type 2 diabetes, is increasingly common among young adults due to rising rates of obesity and sedentary lifestyles. Diabetes accelerates the development of atherosclerosis by promoting inflammation, insulin resistance, and endothelial dysfunction (5). High blood glucose levels can damage the blood vessels and increase the risk of plaque rupture, which is a primary trigger for ACS (6). In addition, diabetes often coexists with other risk factors such as hypertension and dyslipidemia, which further increases the likelihood of developing CAD at an early age (7). According to a study by Rask-Madsen et al. (8), diabetic young adults have a significantly higher risk of ACS compared to their non-diabetic counterparts, and effective management of diabetes is crucial for preventing early cardiovascular events. A family history of CAD is an important risk factor for ACS, particularly when it involves first-degree relatives. Genetic predispositions contribute to early-onset CAD and may involve inherited lipid disorders, early-onset hypertension, or other metabolic conditions that increase the risk of atherosclerosis (9). Individuals with a family history of heart disease are more likely to develop CAD at a younger age, which predisposes them to earlier ACS events. According to the Framingham Heart Study (10), the risk of cardiovascular disease doubles if a first-degree relative experiences a heart attack before the age of 55. Early screening and preventive measures are essential for individuals with a family history of cardiovascular disease to mitigate the risk of ACS. Chronic stress is increasingly recognized as an important modifiable risk factor for ACS in young adults. Stress can trigger the release of stress hormones such as cortisol and adrenaline, which can increase blood pressure, heart rate, and contribute to inflammation, all of which accelerate atherosclerosis and increase the likelihood of ACS (11). This study aimed to assess the prevalence of risk factors of ACS in young adults.

II. Methods

This cross-sectional observational study was conducted at the Department of Cardiology 250 Baded General Hospital, Naogaon, Bangladesh. The study duration was one year. The study population consisted of a total of 100 young adults under the age of 35 years who were admitted to the study hospital for some form of acute coronary syndromes and had undergone coronary angiograms during the study period. Informed written consent was obtained from all the participants before data collection, and ethical approval was obtained from the ethical review committee of the study hospital. Data were collected using the predesigned semi-structured data collection sheet. Data analysis was carried out by using the statistical package SPSS version 23.0 Windows software. Categorical data were analyzed by the Chi-Square test. A p-value of less than 0.05 was considered significant.

Inclusion Criteria

- Patients aged 35 years or younger
- Patients who underwent coronary angiogram
- Patients who had given consent to participate in the study.

Exclusion Criteria

- Patients over the age of 35 years
- Unable to answer the criteria question.
- Exclude those affected with other chronic diseases etc.

III. Results

Table 1: Distribution of the participants by physical characteristics (n=100)

Characteristics	n	%
Gender		
Male	90	90.0
Female	10	10.0
Age		
<20	5	5.0
21-30	28	28.0
31-35	67	67.0
BMI		
Normal (18.0-22.9)	83	83.0
Overweight (23.0-24.0)	5	5.0
Obese (>24.0)	12	12.0

Among the participants (n=100), males constituted the majority (90%), with females accounting for only 10%. The age distribution showed that 5% were aged ≤20 years, 28% were between 21-30 years, and the majority

(67%) were between 31-35 years. Regarding BMI, 83% of participants had normal weight (18.0-22.9), 5% were overweight (23.0-24.0), and 12% were classified as obese (>24.0). [Table 1]

Table 2: Distribution of study participants according to stress levels (n=100)

Stress Level	n	%
No stress	63	63.0
Mild Stress	22	22.0
Moderate Stress	15	15.0

Among the participants (n=100), 63% reported no stress, while 22% experienced mild stress. Moderate stress was observed in 15% of the participants. [Table 2]

Table 3: Distribution of the study subjects according to clinical presentation (n=100)

Clinical presentations	n	%
Chest pain Atypical	5	5.0
Chest pain Typical	95	95.0
Shortness of breath	30	30.0
Nausea/vomiting	10	10.0
Palpitation	8	8.0
Sweating	5	5.0

Chest pain was the most common clinical presentation among participants (n=100), with 95% reporting typical chest pain and 5% reporting atypical chest pain. Shortness of breath was noted in 30%, nausea/vomiting in 10%, palpitations in 8%, and sweating in 5% of the participants. [Table 3]

Table 4: Risk factor distribution of the participants (n=100)

Risk Factors	n	%
Family History of CAD	20	20.0
Smoking	43	43.0
Diabetes Mellitus	55	55.0
Hypertension	70	70.0
Dyslipidemia	48	48.0

Among the participants (n=100), hypertension was the most prevalent risk factor, affecting 70%, followed by diabetes mellitus (55%) and dyslipidemia (48%). Smoking was observed in 43% of the participants, while 20% had a family history of coronary artery disease (CAD). [Table 4]

Table 5: Distribution of the study subjects according to ECG findings (n=100)

ECG findings	n	%
LBBB*	3	3.0
Normal sinus rhythm	8	8.0
Sinus Tachycardia	18	18.0
ST segment depression	10	10.0
ST-segment elevation	50	50.0
T wave changes	13	13.0

*LBBB- Left Bundle Branch Block

ECG findings among participants (n=100) revealed that 50% had ST-segment elevation, while 18% showed sinus tachycardia and 13% exhibited T wave changes. ST segment depression was observed in 10%, normal sinus rhythm in 8%, and left bundle branch block (LBBB) in 3% of the participants. [Table 5]

Table 6: Distribution of the study subjects according to lipid profile (n=100)

Lipid profile	Mean ± SD	Range (min-max)
Triglycerides (mg/dl)	212.0 ± 52.9	91.0 – 368.0
Total cholesterol (mg/dl)	201.8 ± 81.8	66.0 – 446.0
LDL-C (mg/dl)	138.3 ± 46.1	48.0 – 268.0
HDL-C (mg/dl)	39.1 ± 6.1	29.0 – 49.0

The lipid profile analysis of participants (n=100) showed elevated mean triglyceride levels (212.0 ± 52.9 mg/dl) with a range of 91.0 to 368.0 mg/dl. The mean total cholesterol level was 201.8 ± 81.8 mg/dl, ranging from 66.0 to 446.0 mg/dl. LDL-C levels were higher than normal, with a mean of 138.3 ± 46.1 mg/dl (range: 48.0 – 268.0 mg/dl). Meanwhile, HDL-C levels were within normal limits, with a mean of 39.1 ± 6.1 mg/dl (range: 29.0 – 49.0 mg/dl). [Table 6]

IV. Discussion

The age distribution of our study participants reveals that a majority (67%) were between the ages of 31-35, followed by 28% in the 21-30 age group, with only 5% under 20 years of age. These findings are consistent with trends reported in other studies, which suggest that ACS incidence in younger adults is on the rise, particularly in individuals aged 30-40 years (12). The BMI distribution in our cohort indicates that the majority (83%) of participants were classified as having a normal BMI, with a smaller proportion being overweight (5%) or obese (12%). These figures contrast with findings from other studies, where overweight and obesity have been identified as significant risk factors for ACS in younger adults. For instance, a study by Lavie et al. (13) showed that obesity is directly associated with increased cardiovascular risk and adverse outcomes in young adults. Stress levels in our study showed that a majority (63%) of participants reported no stress, with 22% experiencing mild stress and 15% reporting moderate stress. While the majority of participants had low-stress levels, moderate stress, which has been linked to a higher incidence of cardiovascular events, was still prevalent in a notable portion of the cohort. The impact of psychological stress on cardiovascular health, particularly in young adults, has been well-documented in prior research. A study by Steptoe et al. (11) concluded that chronic psychological stress significantly increases the risk of ACS by influencing inflammatory markers and contributing to endothelial dysfunction. The clinical presentations in our study showed that the majority (95%) of participants experienced typical chest pain, which is consistent with the classical presentation of ACS. However, 5% of participants reported atypical chest pain, which could be indicative of less common ACS presentations, especially in younger individuals. Atypical presentations in younger adults are often missed or misdiagnosed, leading to delayed treatment. Our findings are similar to those of a study by McGill et al. (14), which noted that atypical presentations are more common in younger adults with ACS, leading to a higher risk of diagnostic errors and delayed interventions. The risk factor distribution in this study revealed a high prevalence of hypertension (70%), followed by diabetes mellitus (55%), dyslipidemia (48%), and smoking (43%). Hypertension was the most prevalent risk factor, consistent with global studies that highlight its role as a key risk factor for early-onset ACS (15). The association between hypertension and increased cardiovascular risk is well-established, as it promotes atherosclerosis and plaque rupture, which can lead to acute coronary events. Similarly, the high rates of diabetes and dyslipidemia observed in this study align with findings from other studies, which report that metabolic disturbances significantly contribute to the development of early-onset ACS (16). Smoking remains one of the most modifiable risk factors for ACS, and efforts to reduce smoking rates among young adults are critical for preventing early coronary events. Interestingly, only 20% of participants reported a family history of coronary artery disease (CAD), which is lower than what is seen in some populations with a higher genetic predisposition to CAD (10). The lipid profile analysis revealed elevated mean triglyceride levels (212.0 ± 52.9 mg/dl) and LDL-C levels (138.3 ± 46.1 mg/dl), the mean HDL-C levels (39.1 ± 6.1 mg/dl) were within the lower end of the normal range, and low HDL-C is another important contributor to cardiovascular risk. These findings are consistent with studies showing that dyslipidemia, particularly elevated triglycerides and LDL-C, plays a significant role in the development of ACS in younger individuals (17).

Limitations of The Study

The study was conducted in a single hospital with a small sample size. So, the results may not represent the whole community.

V. Conclusion

This study concludes that the increasing prevalence of ACS among young adults emphasizes the role of traditional risk factors such as hypertension, diabetes, dyslipidemia, smoking, and stress. The clinical and diagnostic findings in this study, including typical chest pain, ST-segment elevation on ECG, and elevated lipid levels, align with previous research, underlining the need for early detection and intervention to reduce the burden of ACS in younger populations.

VI. Recommendation

Based on the findings, it is recommended that healthcare professionals focus on early screening and intervention for young adults at risk of acute coronary syndrome (ACS), particularly those with hypertension, diabetes, dyslipidemia, and smoking habits. Lifestyle modifications, including smoking cessation, weight management, and stress reduction, should be emphasized.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee

References

- [1] Lloyd-Jones Dm, Nam Bh, D'agostino Sr Rb, Levy D, Murabito Jm, Wang Tj, Et Al. Parental Cardiovascular Disease As A Risk Factor For Cardiovascular Disease In Middle-Aged Adults: A Prospective Study Of Parents And Offspring. *Jama*. 2004;291(18):2204–11.
- [2] Gooding Hc, Gidding Ss, Moran Ae, Redmond N, Allen Nb, Bacha F, Et Al. Challenges And Opportunities For The Prevention And Treatment Of Cardiovascular Disease Among Young Adults: Report From A National Heart, Lung, And Blood Institute Working Group. *J Am Heart Assoc*. 2020 Oct 6;9(19):E016115.
- [3] Duncan Ms, Freiberg Ms, Greevy Ra, Kundu S, Vasan Rs, Tindle Ha. Association Of Smoking Cessation With Subsequent Risk Of Cardiovascular Disease. *Jama*. 2019;322(7):642–50.
- [4] Arnett Dk, Blumenthal Rs, Albert Ma, Buroker Ab, Goldberger Zd, Hahn Ej, Et Al. 2019 Acc/Aha Guideline On The Primary Prevention Of Cardiovascular Disease: A Report Of The American College Of Cardiology/American Heart Association Task Force On Clinical Practice Guidelines. *Circulation* [Internet]. 2019 Sep 10 [Cited 2024 Nov 20];140(11). Available From: <https://www.ahajournals.org/doi/10.1161/Cir.0000000000000678>
- [5] Pearson Ta, Mensah Ga, Alexander Rw, Anderson JI, Cannon Ro, Criqui M, Et Al. Markers Of Inflammation And Cardiovascular Disease: Application To Clinical And Public Health Practice: A Statement For Healthcare Professionals From The Centers For Disease Control And Prevention And The American Heart Association. *Circulation*. 2003 Jan 28;107(3):499–511.
- [6] Rask-Madsen C, King Gl. Vascular Complications Of Diabetes: Mechanisms Of Injury And Protective Factors. *Cell Metab*. 2013;17(1):20–33.
- [7] Ceriello A, Standl E, Catrinioiu D, Itzhak B, Lalic Nm, Rahelic D, Et Al. Issues Of Cardiovascular Risk Management In People With Diabetes In The Covid-19 Era. *Diabetes Care*. 2020;43(7):1427–32.
- [8] Rask-Madsen C, Kahn Cr. Tissue-Specific Insulin Signaling, Metabolic Syndrome, And Cardiovascular Disease. *Arterioscler Thromb Vasc Biol*. 2012 Sep;32(9):2052–9.
- [9] Rankinen T, Sarzynski Ma, Ghosh S, Bouchard C. Are There Genetic Paths Common To Obesity, Cardiovascular Disease Outcomes, And Cardiovascular Risk Factors? *Circ Res*. 2015 Feb 27;116(5):909–22.
- [10] D'agostino Rb, Vasan Rs, Pencina Mj, Wolf Pa, Cobain M, Massaro Jm, Et Al. General Cardiovascular Risk Profile For Use In Primary Care: The Framingham Heart Study. *Circulation*. 2008 Feb 12;117(6):743–53.
- [11] Steptoe A, Kivimäki M. Stress And Cardiovascular Disease. *Nat Rev Cardiol*. 2012;9(6):360–70.
- [12] Nagaraj N. Study Of Myocardial Infarction In Young Adults [Internet] [Phd Thesis]. Rajiv Gandhi University Of Health Sciences (India); 2010 [Cited 2024 Nov 20]. Available From: <https://search.proquest.com/openview/514b4a31b15330f3d128bd7df92ca116/1?pq-origsite=Gscholar&Cbl=2026366&Diss=Y>
- [13] Cj L. Obesity And Cardiovascular Disease: Risk Factor, Paradox, And Impact Of Weight Loss. *J Am Coll Cardiol*. 2009;53:1925–32.
- [14] McGill Hc, McMahan Ca, Zieske Aw, Tracy Re, Malcom Gt, Herderick Ee, Et Al. Association Of Coronary Heart Disease Risk Factors With Microscopic Qualities Of Coronary Atherosclerosis In Youth. *Circulation*. 2000 Jul 25;102(4):374–9.
- [15] Chobanian Av, Bakris Gl, Black Hr, Cushman Wc, Green La, Izzo JI, Et Al. Seventh Report Of The Joint National Committee On Prevention, Detection, Evaluation, And Treatment Of High Blood Pressure. *Hypertension*. 2003 Dec;42(6):1206–52.
- [16] Arora S, Stouffer Ga, Kucharska-Newton Am, Qamar A, Vaduganathan M, Pandey A, Et Al. Twenty-Year Trends And Sex Differences In Young Adults Hospitalized With Acute Myocardial Infarction: The Aric Community Surveillance Study. *Circulation*. 2019 Feb 19;139(8):1047–56.
- [17] Dogan C, Bayram Z, Karagoz A, Bakal Rb, Erdogan E, Yilmaz F, Et Al. Is Elevated Triglyceride High-Density Lipoprotein Cholesterol Ratio A Risk Factor That Causes Acute Coronary Syndrome To Appear Earlier? *Clin Study*. 2019;770:775.