

Perceived Stigma and Medication Adherence among Mentally ill Patients Visiting Psychiatric OPD of Chitwan Medical College at Bharatpur, Chitwan

Sarita Nepal¹, Amrita Ghimire¹, Ramesh Subba¹, Alisha Joshi¹,

Dr Khagendra Kafle², Dr Sukra Sagar Pandit²

Department of Psychiatric Nursing, Chitwan Medical College, Bharatpur-10

Department of Psychiatry, Chitwan Medical College, Bharatpur 10

Abstract

Background: the perceived or anticipated stigma is associated with patients' beliefs about the attitudes of the general public to the members of stigmatized groups. The aim of this study is to find out level of stigma and medication adherence.

Methods: Descriptive cross sectional research design was performed among mental illness patient visiting on psychiatric OPD of Chitwan Medical College Teaching Hospital. A total of 106 respondents were selected by adopting Non probability Convenience sampling technique. Data was collected by using structured interview schedule. Collected data was analyzed and interpreted through descriptive inferential statistic.

Results: The findings of this study revealed that highest 68.9% respondents was found having mild level of stigma where as 23.6% was found having moderate level of stigma & remaining 7.5% had minimal level of stigma perceived. Similarly lowest 92.5% respondent was found having low level of medication adherence where as 7.5% respondent was found medium level of medication adherence. There is a significant correlation between self-stigma and adherence to treatment.

Conclusion: The study showed a significant correlation between self-stigma and adherence to treatment. High levels of self-stigma are associated with discontinuation of medications without a psychiatrist's recommendation. This connection was present in all diagnostic groups.

Keywords: self-stigma, adherence, discontinuation of drugs, mental disorders

Date of Submission: 23-12-2024

Date of Acceptance: 03-01-2025

I. Introduction

Stigma is a negative differentiation attached to some members of society who are affected by some particular condition or state "The term stigma connotes a deep mark of shame and degradation carried by a person as a function of being a member of a devalued social group." The stigmatized individual experiences social distancing, fear, rejection and ill treatment from others in the society¹ Individuals with mental illness have the dual burden

of coping with the symptoms of mental illness, like hallucinations, depression, delusions and anxiety, as well as the societal stigmatization of their illness.² Stigma is a powerful social process that is characterized by labelling, stereotyping, and separation, leading to status loss and discrimination, all occurring in the context of power.³

Stigma is a basic component of the negative discrimination that people with mental illness experience every day. It blocks access to facilities that have been created to help people with mental illness. Furthermore, the attitudes held by people with authority (including clinical staff and officers) towards people with mental illness are likely to influence their attitude towards them and hence the experience and treatment outcome of the patients⁴

The public stigma on mental illnesses manifests itself in the way the population reacts to mental. Public stigma have negative effects on the lives of people with mental illness, by preventing them from pursuing vocational, housing, and healthcare goals, and holding them back from seeking treatment and affecting the quality of delivered healthcare⁵

Experienced stigma is characterized by the perception of being a victim (recently or throughout life) of stereotypes, prejudice and discrimination. The anticipated stigma relates to the expectation of discrimination (Fox et al., 2017), and they are sometimes associated with internalized stigma (Bos et al., 2013). The

internalized stigma can be defined as the internalization of stereotypes, prejudices and social discrimination to the Self (Corrigan et al., 2006; Fox et al., 2017). These types of MIS are related to delayed treatment seeking and adherence by people with mental illness (PWMI) (Bos et al., 2013; Corrigan et al., 2006; Fox et al., 2017).⁶

Psychiatric stigma research has been, up until recently, primarily focused on public concepts of mental illness and the negative reactions toward mentally ill persons displayed by individuals or societal groups 1–3. Several disease and patient characteristics have been identified as factors influencing stigmatization 4,5, and anti-stigma initiatives have been established to decrease stigmatizing attitudes and discriminating actions in individuals and society as a whole. In recent years, the proposed inclusion of the attenuated psychosis syndrome in the DSM-5 has raised concerns about the potential stigmatization of patients being labelled “at-risk” 13–16, particularly considering the high number of false positives (only approximately 30% of patients showing putative initial prodromal symptoms eventually develop psychosis within the following 2.5 years)⁷

In most societies, mental illness carries a substantial stigma. It is considered as an amalgamation of three related problems: a lack of knowledge (ignorance), negative attitudes (prejudice), and exclusion or avoidance behaviours (discrimination). The mentally ill are labelled as different from other people and are viewed negatively by others. Stigmatization can lower a person’s self-esteem, contribute to disrupted family relationships, and affect employability⁸

II. Methods

Descriptive cross sectional research design was performed among mental illness patient visiting on psychiatric OPD of Chitwan Medical College Teaching Hospital. A total of 106 respondents’ were selected by adopting Non probability Convenience sampling technique. Data was collected by using structured interview schedule. Collected data was analyzed and interpreted through descriptive inferential statistic. The research was done for four weeks periods.

Research approval was obtained from Nursing Research Committee of School of Nursing, Chitwan Medical College (CMC) and ethical approval was obtained from CMC, Institutional Review Committee (IRC), Bharatpur -10, Chitwan. The verbal informed consent was obtained from each respondent by clarifying the purpose of the study prior to data collection. Each respondent was allowed to have a voluntary choice for participation. Respondent’s dignity was maintained by giving right to reject or discontinue from the research study at any time. Confidentiality of the information was maintained by not disclosing the information and using the information only for the research purpose.

III. Results

Out of 106 respondents, maximum respondents 67% belonged to age group more or equal to 43 years. Among them 52.8% were male. Similarly 80.2% belong in Hindu religion. Whereas highest 36.8% had basic education. More than half 68.9% of respondents were married similarly 91.5% respondents had family support and 91.5% had sufficient income source.(Table1)

TABLE 1
Respondents’ Socio-demographic characteristics
n=106

Variables	Frequency	percentage
Age		
<43.27	71	67
≥43.27	35	33
Mean ±SD= 43.27±8.169, Min=25,Max=60		
Gender		
Male	56	52.8
Female	50	47.2
Religion		
Hindu	85	80.2
Non Hindu	21	19.8
Education		
Illiterate	22	20.8
Can read and write	29	27.4
Basic	39	36.8
Secondary	8	7.5
Bachelor and above	8	7.5
Occupation		
Home maker	51	48.1

Government job	9	8.5
Self employment	10	9.4
Others	36	34
Marital status		
Married	73	68.9
Unmarried	13	12.3
Divorce	12	11.3
Separate	8	7.5
Family type		
Nuclear	69	65.1
Joint	29	27.4
Extended	8	7.5
Family support		
Yes	97	91.5
No	9	8.5
Monthly income		
Sufficient	97	91.5
Insufficient	9	8.5

highest 35.84% respondents had diagnosis of depressive with psychotic similarly 66% respondents had less than 5 years duration of onset of disorder likewise 55.66% had need of hospitalization .Highest 66% had more than one week hospitalization and 57.54% respondents length of treatment was more than 2 years.(Table 2)

TABLE 2
Respondent's disease related characteristic
n=106

Variables	Frequency	percentages
Diagnosis		
Depression with psychotic	38	35.84
Depression without psychotic	17	16%
Schizophrenia	27	25.47%
BPAD	15	14.15
Psychosis	9	8.4%
Duration		
<5 years	70	66%
>5years	36	33.95%
Hospitalization		
Yes	59	55.66%
No	47	44.33%
If yes, then stay of last hospitalization (n=59)		
<1 week	20	33%
>1 week	39	66%
Length of medicine use		
<2 years	45	42.45%
>2 years	61	57.54%
Side effect of medicine		
Yes	36	33.95%
No	70	66%
If yes, types of effect (n=36)		
Nausea	18	50%
Vomiting	10	27.77%
Headache	3	8.33%
Drowsy	5	13.88%

TABLE 3

Respondent Response Regarding IMCI Scale Based on different Domain
TABLE 3. A Alienation Related Statement

n=106

Domain 1: Alienation	Strongly disagree	Disagree	Agree	Strongly agree
I feel out of place in the world because I have mental illness	8 (7.5%)	26 (24.5)	55 (51.9)	17 (16%)
I am embarrassed or ashamed that I have a mental illness	17 (16%)	44 (41.5)	45 (42.5)	0
I feel inferior to others who do not have a mental illness	25 (23.6%)	56 (52.8%)	25 (23.6%)	0
I am disappointed in myself for having a mental illness	8 (7.5%)	52 (49.1%)	46 (43.4%)	0
Having a mental illness has spoiled my life	25 (23.6%)	30 (28.3%)	51 (48.1%)	0
People without mental illness could not possibly understand me	8 (7.5%)	72 (67.9%)	18 (17%)	8 (7.5%)

TABLE3.B Stereotype Endorsement Related Statement

n=106

Domain 2: Stereotype	Strongly disagree	Disagree	Agree	Strongly agree
Mentally ill people tend to be violent	8 (7.5%)	77 (72.6%)	21 (19.8%)	0
Mentally ill people shouldn't get married	25 (23.6%)	77 (72.6%)	4 (3.8%)	0
People with mental illness cannot live a good ,rewarding life	16 (15.1%)	44 (41.5%)	46 (43.4%)	0
People can tell that i have a mental illness by the way i look	8 (7.5%)	38 (35.8%)	60 (56.6%)	0
Because i have a mental illness ,i need other to make most decisions for me	16 (15.1%)	21 (19.8%)	69 (65.1%)	0
I can't contribute anything to society because i have a mental illness	33 (31.1%)	59 (55.7%)	14 (13.2%)	0
Stereotype about the mentally ill apply to me	15 (14.1%)	53 (50%)	35 (33%)	3 (2.8%)

TABLE 3.C Perceived Discrimination Related Statement

Domain 3: Perceived Dricrimination	Strongly disagree	Disagree	Agree	Strongly Agree
People discriminate against me because I have a mental illness	6 (5.6%)	35 (33%)	59 (55.6%)	6 (5.6%)
Being around people who do not have a mental illness makes me feel out of place or inadequate	8 (7.5%)	64 (60,4%)	34 (32.1%)	0
I stay away from social situations in order to protect my family or friends from embarrassment	15 (14.1%)	50 (47.1%)	39 (36.8%)	2 (1.8%)

TABLE 3.D Social Withdrawal Related Statement

Domain 4 : Social Withdrawal	Strongly disagree	Disagree	Agree	Strongly agree
I avoid getting close to people who don't have a mental illness to avoid rejection.	8 (7.5%)	80 (75.4%)	17 (16%)	1 (0.9%)
I don't socialize as much as I used to because my mental illness might make me look or behave "weird."	25 (23.6%)	39 (36.8%)	42 (39.6%)	0
I don't talk about myself much because I don't want to burden others with my mental illness.	8 (7.5%)	48 (45.3%)	50 (47.2%)	0
Negative stereotypes about mental illness keep me isolated from the "normal" world.	8 (7.5%)	48 (45.3%)	45 (42.4%)	5 (4.6%)
People often patronize me, or treat me like a child, just because I have a mental illness.	9 (8.4%)	52 (49%)	45 (42.4%)	0
People ignore me or take me less seriously just because I have a mental illness.	8 (7.5%)	39 (36.8%)	54 (50.9%)	5 (4.6%)
Nobody would be interested in getting close to me because I have a mental illness.	12 (11.3%)	27 (25.5%)	65 (61.3%)	2 (1.8%)

Others think that I can't achieve much in life because I have a mental illness.	8 (7.5%)	12 (11.3%)	64 (60.4%)	22 (20.8%)
---	-------------	---------------	---------------	---------------

TABLE 3.E Stigma Resistance

Domain5: Stigma resistance	Strongly disagree	Disagree	Agree	Strongly agree
People with mental illness make important contributions to society	30 (28.3%)	50 (47.1%)	16 (15%)	6 (5.6%)
I feel comfortable being seen in public with an obviously mentally ill person	60 (56.6%)	35 (33%)	8 (7.5%)	3 (2.8%)
Living with mental illness has made me a tough survivor	8 (7.5%)	61 (57.5%)	29 (27.4%)	8 (7.5%)
In general, I am able to live my life the way I want to	8 (7.5%)	55 (51.9%)	27 (25.5%)	16 (15%)
I can have a good, fulfilling life, despite my mental illness	33 (31.1%)	44 (41.5%)	27 (25.5%)	2 (1.8%)

**TABLE 4
Mean Score of Perceived Stigma Among Mental Illness Patient
n=106**

Domain	No items	Maximum Possible Score	Obtained Score range	Mean ±SD	Mean %
Alienation	6	24	8-19	13.88±2.79	56.83
Stereotype endorsement	7	28	7-19	15.27±3.17	54.31
Perceived discrimination	3	12	3-9	6.95±1.49	56.19
Social withdrawal	8	32	8-24	19.37±3.94	60.53
Stigma resistance	5	20	10-19	12.96±2.43	64.85
Total	29	116	36-90	68.43±13.82	58.54

Table 4 shows Mean Score of perceived stigma among mentally ill patient on five different domains. Out of 106 respondents, highest number 64.85% respondents had stigma resistance and 54.31% respondents had stereotype endorsement

**TABLE 5
Level of Stigma Perceived By Respondents'
n=106**

Level of stigma	Frequency	Percentage (%)
Minimal	8	7.5
Mild	73	68.9
Moderate	25	23.6

Minimal (1.00-2.00), Mild (2.01-2.50), Moderate (2.51-3.00)

Table 5 shows highest 68.9% respondents was found having mild level of stigma where as 23.6% was found having moderate level of stigma & remaining 7.5% had minimal level of stigma perceived.

**TABLE 6
Level of medication adherence**

Level of medication adherence	Frequency	Percentage (%)
Low	98	92.5
Medium	8	7.5

Table 6 shows lowest 92.5% respondent was found having low level of medication adherence where as 7.5% respondent was found medium level of medication adherence.

IV. Discussion

Descriptive cross sectional research design was used to find out perceived stigma and medication adherence among mental illness patient. The study population consisted of 106 attended at psychiatric OPD at Chitwan Medical College. Highest 68.9% respondents was found having mild level of stigma where as 23.6%

was found having moderate level of stigma & remaining 7.5% had minimal level of stigma perceived. In this study lowest 92.5% respondent was found having low level of medication adherence where as 7.5% respondent was found medium level of medication adherence. The study showed a significant correlation between self-stigma and adherence to treatment. Medication adherence was associated with lower perceived stigma.

This study supported on the topic of Perceived Self-Stigma, Adherence to Treatment and Discontinuation of Medication among Schizophrenic Patients on Cairo, Egypt. Cairo, Egypt results revealed that there is a positive association between stigma and discontinuation of medications. The conclude the level of internalized stigma of patients with schizophrenia was determined to be high and the internalized stigma had a negative impact on the adherence and attitude toward the treatment.⁹

This study supported by the study conducted on outpatients attending the Department of Psychiatry, University Hospital Olomouc, Czech Republic on 332 patients. The level of self-stigma was measured with the Internalized Stigma of Mental Illness and adherence with the Drug Attitude Inventory. Showed a significant correlation between self-stigma and adherence to treatment. High levels of self-stigma are associated with discontinuation of medications without a psychiatrist's recommendation. This connection was present in all diagnostic groups.¹⁰

The study revealed that there was high self-stigma among patients with mental illness and a significant association between overall ISMI score and level of medication adherence. These require mental health professionals. number of relapses (std. $\beta = 0.183$, $p < 0.01$) and medication nonadherence (std. $\beta = 0.084$, $p = 0.021$) were positively associated with self-stigma¹¹

V. Conclusion

It is concluded that more than half of respondent have minimal level of stigma and Similarly maximum respondent was found having low level of medication adherence. There is a significant correlation between self-stigma and adherence

References:

- [1] Dharitri R, Rao Sn, S Kalyanasundaram. Stigma Of Mental Illness: An Interventional Study Reduces Its Impact In The Community: Indian J Psychiatry. 2015 Apr-Jun; 57(2): 165–173
- [2] Abdisa E, Fekadu G, Girma S Etl. Self Stigma And Medication Adherence Among Patients With Mental Illness Treated At Jimma University Medical Center, Southwest Ethiopia: Abdisa Et Al. Int J Ment Health Syst (2020) 14:56
- [3] Nyblade L, Stockton M A, Giger K Etl. Stigma In Health Facilities: Why It Matters And How We Can Change It: Bmc Medicine Volume 17, Article Number: 25 (2019)
- [4] Eissa Ab, Elhabiby M, Serafi De. Investigating Stigma Attitudes Towards People With Mental Illness Among Residents And House Officers: An Egyptian Study. Middle East Current Psychiatry. Volume 27 Article Number: 18 (2020)
- [5] Yin H, Wardenaar KJ, Xu G, Tain H & Schoevers Ra. Mental Health Stigma And Mental Health Knowledge In Chinese Population: A Cross-Sectional Study: Bmc Psychiatry Volume 20, Article Number: 323 (2020)
- [6] Fox A, Earnshaw Va, Taverna E, Vogt A. Conceptualizing And Measuring Mental Illness Stigma: The Mental Illness Stigma Framework And Critical Review Of Measures: 2017 (3) 4
- [7] Gerlinger G, Marta H \$ Christoph U. Personal Stigma In Schizophrenia Spectrum Disorders: A Systematic Review Of Prevalence Rates, Correlates, Impact And Interventions. World Psychiatric 2013 June; 12(2):155-164
- [8] Eissa Am, Mahmoud M, Doha Ei Etl. Investigating Stigma Attitudes Towards People With Mental Illness Among Residents And House Officers: An Egyptian Study: Middle East Current Psychiatry Volume 27, Article Number: 18 (2020)
- [9] Kamaradova D, Latalova E. Connection Between Self-Stigma, Adherence To Treatment, And Discontinuation Of Medication: Patient Preference And Adherence Volume 10(Issue 1):1289-1298
- [10] Asrat B, Eskeziya A A & Yimer T. Internalized Stigma Among Patients With Mental Illness Attending Psychiatric Follow-Up At Dilla University Referral Hospital, Southern Ethiopia: Volume 2018 |Article Id 1987581
- [11] Abdisa E, Fekadu G, Grima S. Self Stigma And Medication Adherence Among Patients With Mental Illness Treated At Jimma University Medical Center, Southwest Ethiopia