

## A Prospective Study of Outcome of Pregnancy Complicated By Threatened Abortion

Dr. Manisha Bahad<sup>1</sup>, Meena Bhanwar Singh<sup>2</sup>, Mahila Chikitsalaya<sup>3</sup>

<sup>1</sup>M.S. (Obst & Gynae ) Correspondence Author

<sup>2</sup>Professor and Unit Head

<sup>3</sup>Institution Department of Obstetrics and Gynaecology

---

### Abstract:

**Objective:** To evaluate prospectively the risk of adverse pregnancy outcome in women presenting with threatened miscarriage.

**Method:** A prospective study was performed on 70 women presenting with bleeding in first and early second trimester and 70 asymptomatic age matched controls. Main outcome measures included gestational age and weight at delivery and incidence of adverse pregnancy outcome.

**Results:** The first trimester miscarriage rate, after confirmation of viability in the threatened miscarriage group, was 14.28%. Women presenting with threatened miscarriage were more likely to have PPRM than controls 31.42% compared with 11.42%, respectively ( relative risk 3.5, confidence interval 1.5-8.6 ) They were also more likely to have preterm delivery, 18.57% compared with 5.7%, respectively (relative risk 3.8, CI 1.16-12.2 ); LBW, 25.71% compared with 12.85%, respectively ; antepartum haemorrhage , 7.14% compared with 1.43% and IUGR 7.14% compared with 2.85% respectively. Birth weight 2458 ± 440 g compared with 2665± 500 respectively, gestational age 32.9 ± 9.5 compared with 38.4 ± 6.76 respectively.

**Conclusion:** First-trimester vaginal bleeding is an independent risk factor for adverse obstetric outcome and this risk factor should be taken into consideration when deciding upon antenatal surveillance and management of their pregnancies.

---

### I. Introduction

Threatened miscarriage, defined as vaginal bleeding before 20 weeks of gestation, is a common complication affecting about 20% of pregnancies. It has been shown to be associated with an increased risk of poor obstetric outcomes such as preterm labor, low birth weight, and premature rupture of membranes. Moreover, when pregnant women have bleeding, it may cause stress and anxiety for the mother-to-be about the outcome of pregnancy. So, it is necessary to be diagnosed and managed to prevent maternal or fetal mortalities and morbidities (1).

The scientific literature regarding threatened abortion is relatively limited on the subject of outcomes and viability at term. Small number of patients and significantly biased data collection have limited previous studies of pregnancies that were complicated by threatened abortion (2-4). Many studies suggest that first-trimester vaginal bleeding is associated with a worse outcome (2-9). However, there have been few studies that evaluated outcomes other than viability at term, after the documentation of a living embryo. In general, the incidence of spontaneous abortion after first-trimester bleeding is quoted to be 50% before sonographic evaluation for fetal viability (5, 6). If a viable fetus is noted at ultrasound examination after first-trimester vaginal bleeding, 95% to 98% of such pregnancies will still continue beyond 20 weeks of gestation (5).

The diagnosis of threatened miscarriage is frequently made in clinical practice as a result of taking a history of vaginal spotting and the finding of a closed cervix at subsequent vaginal examination. A definitive diagnosis of threatened miscarriage should be made following ultrasonographic examination, confirming the presence of fetal heart activity in an intrauterine pregnancy (8).

Because bleeding originates from the placenta in most cases of threatened abortion, we hypothesized that pregnancies complicated by a first-trimester threatened abortion could be at increased risk for other "placental related" complications. To evaluate this risk of first-trimester bleeding, we evaluated a variety of outcomes that included pregnancy loss, preterm delivery, intrauterine growth restriction (IUGR), preeclampsia, preterm premature rupture of membranes (PPROM), placental abruption, and Cesarean delivery (9).

First-trimester vaginal bleeding is common, occurring in 15-20% of all viable pregnancies. Despite its common occurrence, the risk of adverse outcome for pregnancies with first-trimester threatened abortion and a living embryo has been defined incompletely (10). However, unexplained prenatal haemorrhage, similar to threatened miscarriage, probably exerts an indirect effect (as they were not independent risk factors for early neonatal death) on perinatal outcome through the increased risk of preterm delivery. This suggests that a

threatened miscarriage is possibly related to uteroplacental dysfunction. The current study was conducted on patients with first-trimester threatened abortion to evaluate the outcome.

## **II. Method**

A prospective case-control study was performed in 70 women presenting with bleeding in the first & early second trimester and 70 age-matched control subjects who attended for routine dating or delivery at Mahila Chikitsalaya of SMS medical College, Jaipur from march 2013 to august 2014..

Baseline data were recorded by questionnaire and patient interview. Subjects were divided into two groups: 1) no bleeding, 2) bleeding (defined as spotting or as similar to menses). Post delivery follow-up was performed by telephone interview or medical record review by the research coordinator at each site. Only primigravida included in study to exclude confounding factors of gravidity and recurrent abortion.

Any incident of preeclampsia, IUGR, intrauterine fetal distress (IUFD), low birth weight (LBW), low lying placenta, placenta previa or low lying placenta, birth weight and neonatal sex were recorded. The following pregnancy outcomes between the two groups were compared: abortion, preterm delivery, IUGR, IUFD, PPRM, LBW, preeclampsia, type of delivery.

The following adverse pregnancy outcomes among the two groups were then compared: IUGR (estimated fetal weight by ultrasound examination of <10th percentile or birth weight of <10th percentile for gestational age), gestational hypertension (blood pressure >140/90 mm Hg on at least two occasions >6 hours apart without evidence of chronic hypertension), preeclampsia (criteria for gestational hypertension and significant proteinuria), preterm labor (labor <37 weeks of gestation), PPRM (membrane rupture <37 weeks of gestation), placental abruption (premature separation of a normally implanted placenta), placenta previa (placenta completely or partially covering the internal os), low lying placenta (placenta edge does not reach the internal os but is in close proximity to it ) and caesarean delivery.

The outcomes were established and analysed with SPSS 13. Univariate and multivariate logistic regression analysis were used to evaluate the association among the two groups with regards to specific pregnancy outcomes.

## **III. Results**

The demographic characteristics of the two groups are summarized in Table 1. Statistically non significant differences were noted among the groups for age and parity .

All subjects who were included in this investigation had a viable pregnancy confirmed by ultrasound examination at the time of trial enrolment. Compared with the control group, patients with vaginal bleeding were significantly more likely to have a spontaneous loss (14.28%).

A statistically significant difference was noted in mean birth weight among the two groups: control, 2665 g  $\pm$  500; case group, 2458  $\pm$  440 g ( $P < 0.001$ ). The mean gestational age at delivery for patients in control group and vaginal bleeding group was 36.4  $\pm$  6.76 weeks and 32.9  $\pm$  9.5 weeks, respectively ( $P = 0.014$ ).

The obstetric outcomes for patients with threatened miscarriage compared with patients without bleeding are described in Table 2. No significant difference in the incidence of IUGR ( $P = 0.24$ ), preeclampsia ( $P = 0.75$ ), type of delivery ( $P = 0.08$ ), IUFD ( $P = 1$ ) or antepartum haemorrhage between the control group and subjects with vaginal spotting was noted.

Statistically significant differences were noted in these complications: preterm delivery, 5.7% in controls compared with 18.57% in case group ( $P = 0.02$ , relative risk 3.8, 95% confidence interval 1.16- 12.2); PPRM, 11.42% compared with 31.42%, ( $P < 0.001$ , relative risk 3.5, 95% confidence interval 1.5- 8.6); LBW, 12.85% compared with 25.71% ( $P = 0.054$ ) respectively, were significantly more common in patients with vaginal bleeding compared with control patients.

## **IV. Discussion**

These data show that threatened miscarriage is not only associated with miscarriage but also with adverse pregnancy outcome. Results from this study confirm findings from other authors, that threatened abortion is associated with an increased risk of certain pregnancy-related complications, namely preterm labour, delivery of low birth weight infants and PPRM (12,13). Generally, high incidence of abortion and complications in threatened miscarriage indicate the necessity of proper programming in care and also educating high risk women. A potential limitation of this study is that the severity of vaginal bleeding was based on a subjective description by the patient. However, the ultimate assessment of vaginal bleeding is based on patient report. Therefore, we believe the results of this study can be applied to clinical practice.

In one study, an increased risk of preeclampsia, preterm delivery, placental abruption, and Cesarean delivery was observed for patients who reported light bleeding, although these risks were low with an OR of < 2.0 (13). For patients who reported heavy vaginal bleeding during the first trimester, they observed increased risks of IUGR, preterm delivery, PPRM placental abruption, and Cesarean delivery. With the exception of

Cesarean delivery (OR, <2.0) these associations appear to be both statistically and clinically significant (13). In our study increased risk of preterm delivery (rr= 3.8), PPRM (rr= 3.5) was occurred.

In 1993, Verma *et al.* reported that pregnancy- induced hypertension was significantly more common in subjects with threatened abortion and a viable pregnancy compared with subjects without vaginal bleeding (6% vs. 4.7%, respectively;  $P < 0.05$ ) (2). Another study did not find an association between first-trimester vaginal bleeding and gestational hypertension but did find that patients with light bleeding were statistically more likely to have preeclampsia (13). This association carried a low OR of < 2.0. Similarly, we don't find significant association between vaginal bleeding and preeclampsia.

Preterm delivery before 37 weeks gestation occurs in 7-11% of pregnancies, but is responsible for 85% deaths of normally formed infants. Despite significant advances in perinatal medicine, the incidence of preterm delivery has remained unchanged. The prediction of preterm delivery from currently available methods is unreliable, therefore, associated risk factors remain an important measure of identifying at-risk pregnancies. This is especially evident by the significantly increased rate of early neonatal death rate in infants born from mother who had presented with a threatened miscarriage, probably secondary to the increased rate of preterm deliveries and placental abruption. The only potential risk factor found to be associated significantly with the risk of preterm delivery in women with a threatened miscarriage was unexplained antepartum haemorrhage rather than other factors, such as smoking or preterm rupture of membranes. In our study, preterm delivery was more common in bleeding group (rr= 3.8).

The association between vaginal bleeding and preterm delivery has also been noted by others (21- 23). Both Batzofin *et al.* (4) and Williams *et al.* (7) reported that patients with bleeding had double the risk of preterm delivery compared with patients without bleeding. The study of Williams *et al.* was limited to first trimester bleeding (7); Batzofin *et al.* included patients with bleeding up to 20 weeks (4). Strobino and Pantel-Silverman failed to show an association between preterm delivery before 36 weeks of gestation with light vaginal bleeding in the first or second trimester of pregnancy . Another study found that preterm delivery is increased significantly in patients with either light (OR, <2.0) or heavy (OR, 3.0) first-trimester bleeding (13).

Other studies were reported that patients with first-trimester threatened abortion are also at increased risk for placental abruption and IUGR (1). Placental haemorrhage may recur later in pregnancy, which results in placental abruption. In present study placental abruption was not significantly more common in case group but no increase risk for IUGR was found.

Haddow *et al.* reported an increased risk for low birth weight (<2500 g) in pregnancies that were complicated by vaginal bleeding (26). Infants of patients with heavy bleeding had nearly a 200 g difference in birth weight compared with control infants after accounting for preterm delivery. In present study risk of LBW was increased .

Our findings corroborate other studies that suggested an association between threatened abortion and PPRM (4-6). Although the cause is unclear, it is hypothesized that disruption of the chorionic-amniotic plane by adjacent haemorrhage may make the membranes more susceptible to rupture (4). Alternatively, the prolonged presence of blood may act as a nidus for intrauterine infection. Persistent or recurrent placental haemorrhage could also stimulate subclinical uterine contractions that result in cervical change and eventual ruptured membranes. In our study PPRM was increased in case group (rr = 3.5).

Currently, there is no information in the literature regarding threatened abortion and Cesarean delivery. One study suggested that a statistical association is present with threatened abortion and risk for cesarean delivery (4). In present study, Caesarean delivery was not higher in case group ( $P = 0.08$ ).

In conclusion, the current study reports that patients with first-trimester threatened abortion are at increased risk for spontaneous loss and adverse pregnancy outcome. For patients who reported vaginal bleeding during the first trimester, we observed increased risks of LBW, preterm delivery, PPRM, placental abruption, and low lying placenta. These associations appear to be both statistically and clinically significant. Because the overall prognosis is favourable, these results can be used to help reassure patients with threatened abortion during the first trimester. At the same time, physicians should be aware of the adverse outcomes that are associated with first-trimester bleeding and remain alert for signs of these complications.

## References

- [1]. Mulik V, Bethel J, Bhal K. A retrospective population- based study of primigravid women on the potential effect of threatened miscarriage on obstetric outcome. *J Obstet Gynaecol.* 2004 Apr;24(3):249-253.
- [2]. Verma SK, Premi HK, Gupta TV, Thakur S, Gupta KB, Randhawa I. Perinatal outcome of pregnancies complicated by threatened abortion. *J Indian Med Assoc.* 1994 Nov;92(11):364-365.
- [3]. Hertz JB, Heisterberg L. The outcome of pregnancy after threatened abortion. *Acta Obstet Gynecol Scand.* 1985;64(2):151-156.
- [4]. Batzofin JH, Fielding WL, Friedman EA. Effect of vaginal bleeding in early pregnancy on outcome. *Obstet Gynecol.* 1984 Apr;63(4):515-518.
- [5]. Farrell T, Owen P. The significance of extrachorionic membrane separation in threatened miscarriage. *Br J Obstet Gynaecol.* 1996 Sep;103(9):926-928.
- [6]. Chung TK, Sahota DS, Lau TK, Mongelli JM, Spencer JA, Haines CJ. Threatened abortion: prediction of viability based on signs and symptoms. *Aust N Z J Obstet Gynaecol.* 1999 Nov;39(4):443-447.

- [7]. Williams MA, Mittendorf R, Lieberman E, Monson RR. Adverse infant outcomes associated with first- trimester vaginal bleeding. *Obstet Gynecol.* 1991 Jul; 78(1):14-18.
- [8]. Park IY, Park CH, Lee G, Shin JC. Prognosis of threatened abortion by embryonic/fetal heart beat rate. *Ultrasound Med Biol.* 2006; 32(5): 264.
- [9]. Bowe P, Murphy H. Complications of pregnancy following threatened abortion. *Ir J Med Sci.* 1987 Nov;156(11):328-329
- [10]. Nagy S, Bush M, Stone J, Lapinski RH, Gardó S. Clinical significance of subchorionic and retroplacental hematomas detected in the first trimester of pregnancy. *Obstet Gynecol.* 2003 Jul;102(1):94- 100.
- [11]. Tannirandorn Y, Sangsawang S, Manotaya S, Uerpairojkit B, Samritpradit P, Charoenvidhya D. Fetal loss in threatened abortion after embryonic/fetal heart activity. *Int J Gynaecol Obstet.* 2003 Jun;81(3):263- 266.
- [12]. Das AG, Gopalan S, Dhaliwal LK. Fetal growth and perinatal outcome of pregnancies continuing after threatened abortion. *Aust N Z J Obstet Gynaecol.* 1996 May;36(2):135-139.
- [13]. Weiss JL, Malone FD, Vidaver J, Ball RH, Nyberg DA, Comstock CH, Hankins GD, Berkowitz RL, Gross SJ, Dugoff L, Timor-Tritsch IE, D'Alton ME; FASTER Consortium. Threatened abortion: A risk factor for poor pregnancy outcome, a population-based screening study. *Am J Obstet Gynecol.* 2004 Mar;190(3):745- 750.

**Table 1** Characteristics & Outcomes in women with threatened miscarriage compared with Asymptomatic controls

Characteristics	Cases (n = 70)	Controls (n=70)	P value
Mean Age(yrs)	22.74+ 2.85	23.21 + 2.46	0.254
Literacy	67.1%	72.9%	0.460
Regular ANC Visits	51.4%	62.9%	0.17
Mean Value of Gestational age at presentation(week)	9.5 + 3.53	11.4 + 3.57	0.001
Mean Gestational age at Birth(week)	32.9 + 9.5	36.4 + 6.76	0.014
Birth weight (gm)	2458 + 440	2665 + 0.50	0.016

**Table 2** Adverse Pregnancy Outcomes in women with threatened miscarriage compared with Asymptomatic controls

Outcome	Case ( n = 70)	Control (n = 70)	P value
IUGR	5(7.14%)	2(2.85%)	0.24
LBW	18(25.71%)	9(12.85%)	0.054
Preterm delivery	13(18.57%)	4(5.7%)	0.02
IUFD	2(2.85%)	2(2.85%)	1
Preeclampsia	5(7.14%)	6(8.57%)	0.75
PPROM	22(31.42%)	8(11.42%)	0.004
APH	5(7.14%)	1(1.43%)	0.09
Caesarean del	22(31.4%)	13(18.6%)	0.08