

Study of Serum C-Reactive Protein In Acute Ischemic Stroke And Its Impact on Stroke Severity And Prognosis.

Dr. Chandrabhushan Sharma¹, Dr. Govind Madhaw²

¹(Associate Professor /Department of medicine/ R.I.M.S/Ranchi University, India)

²(Junior Resident /Department of Medicine/ R.I.M.S/ Ranchi University, India)

Abstract:

Introduction: Acute ischemic stroke (AIS) has several risk factors; atherosclerosis of vessels is the most important one. Atherosclerosis being an inflammatory condition is associated with raised serum level of CRP. Hence the possible role of serum CRP in pathogenesis of AIS is being studied. Many studies support that serum CRP is a marker of stroke severity and prognosis.

Material and methods: The present study 'Serum C-reactive protein in Acute Ischemic stroke and its impact on stroke severity and prognosis' was carried out at Department of medicine, RIMS Ranchi, India during August 2015 to August 2016. Total 50 neuroimaging confirmed cases of AIS were selected randomly for this non-interventional prospective study. Serum CRP was measured in all patients and NIHSS was used for assessing stroke severity.

Results: The mean CRP level was 21.93 ± 3.60 mg/l and only 18% of cases were having normal CRP level. 60% of cases were having serum CRP level in the range of 3-30mg/L and 18% were having even the higher level. Age of the patient significantly correlated with the CRP level ($r=0.294$ at $p=0.05$) but the sex didn't affect the rise in CRP ($p=0.490$). Serum CRP level was affecting the stroke severity significantly ($r=0.643$, $p=0.000$). The mean CRP level in discharged patients was 10.51 ± 13.21 mg/l whereas it was 44.11 ± 29.7 mg/l in those who died in the hospital. Thus CRP level affected mortality of acute ischemic stroke significantly. ($p=0.00$).

Conclusion: CRP is raised in most of the patients of acute ischemic stroke and its elevation reveals the underlying inflammation causing atherosclerosis as well as CRP as a marker of brain injury. Serum C-reactive protein elevation is associated with more severe stroke and poor prognosis.

Keywords: Acute ischemic stroke, C-reactive protein, prognostic factors

I. Introduction

Acute ischemic stroke (AIS) is characterized by the sudden loss of blood circulation to an area of the brain, typically in a vascular territory, resulting in a corresponding loss of neurologic function. Among various risk factors atherosclerosis is the most important. Many studies confirmed that early detection and prevention of atherosclerosis is very effective in decreasing the risk of acute ischemic stroke. Atherosclerosis is considered a chronic inflammatory response by vascular endothelium.^[2] C-reactive protein (CRP) is an acute phase reactant produced in response to inflammatory process and therefore it is considered a sensitive marker of inflammation and atherosclerosis.

Cerebrovascular ischemia has been associated with bronchial and periodontal infections.^[3] As a marker of infection and inflammation, high CRP has been associated with acute stroke. Several studies have assessed the value of CRP in the early phase of stroke as a prognostic factor of functional outcome. In one study, patients with CRP levels ≥ 7 mg/L had higher NIHSS scores on admission.^[4] Patients with CRP levels ≥ 7 mg/L taken as a group had worse outcomes in terms of morbidity and mortality.^[5] Interestingly, recent experimental studies have shown that CRP itself may contribute to secondary brain damage after focal cerebral ischemia, possibly via a complement-mediated exacerbation of tissue injury. Since infectious and inflammatory conditions are much prevalent in India compared to developed countries, and only a few studies are available from India on the association of CRP with ischemic stroke, aim of the present study was to assess the possible relationship of CRP in acute ischemic stroke and its impact on stroke severity and mortality.

II. Material and Method

Source of data: Rajendra institute of medical sciences, Ranchi

Study Design: observational and prospective study design.

Duration of study: August 2015 to August 2016.

Ethical consideration: Prior approval for this study was granted by the 'institutional Ethics Committee, RIMS Ranchi.' Written informed consent was obtained from the subject or from their relatives.

Sample Size: 50 patients.

Diagnosis of stroke was confirmed by Brain imaging: CT-scan Or MRI.

Venous blood samples for the estimation of C-reactive protein(CRP) and other biochemical parameters were collected within 72 hours of admission. Within one hour of collection, the samples were centrifuged to separate the serum and were kept in -70°C . CRP was measured using immunoturbidimetric assays (VITROS 5.1). As per the normative data from VITROS manual and current literature, the cardiovascular risk was determined as low risk with CRP levels < 1.0 mg/L, medium risk if $1.0-3.0$ mg/L, high risk when > 3.0 mg/L.^[6]For our study we considered CRP level of ≥ 3 mg/L as high risk and ≤ 3 mg/L as low risk. Detailed information about past medical and surgical history (in-particular infections, systemic inflammation) and other risk factors of cardiovascular and Cerebro-Vascular accident was collected from all patients. Erythrocyte sedimentation rate (ESR) was measured in all subjects. Other investigation included Serum total cholesterol, blood sugar, ECG, RFT etc. The severity of stroke was measured on admission by the *NIH stroke severity scale*. The NIHSS is composed of 11 items, each of which scores a specific ability between 0 and 4. For each item, a score of 0 typically indicates normal function in that specific ability, while a higher score is indicative of some level of impairment. The individual scores from each item are summed to calculate a patient's total NIHSS score. The maximum possible score is 42, with the minimum score being a 0. Score <4 is having Minor stroke, 5-15 for moderate severity, 16-20 for moderate to severe stroke and 21-42 score shows very severe stroke. Patient's outcome/prognosis was recorded as either death in hospital or return home with partial/full recovery.

Inclusion criteria:

- All the patients of acute ischemic stroke confirmed by cerebral imaging, admitted in the department of medicine, RIMS, Ranchi.
- Presented within 72 hours of the onset of stroke.

Exclusion criteria:

- Age < 14 years
- Impaired renal function
- Elevated ESR value
- Severe sepsis, peritonitis, pancreatitis and any other systemic inflammatory conditions
- Patient unwilling to give informed consent.

III. Statistical Analysis

Categorical variables were analyzed using either the chi-square statistic or Fischer exact test, as appropriate. Continuous variables were expressed as mean and statistical significance was tested using Student's t-test. Patients were grouped as normal versus elevated based upon serum admission CRP levels. All other parameters like patient characteristics and risk factors as well as presenting symptoms were dichotomized into 'yes=1' or 'no=0' to make the analysis easy. Also the patient outcome was dichotomized into 'death=1' or 'discharged=0'. Multivariate logistic regression models were constructed to determine variables that predicted abnormal elevations in admission serum CRP. Similar models were constructed to identify variables that independently predicted long-term outcomes. Associations were considered significant if p value is less than 0.05. All analyses were performed with SPSS v20.0.

IV. Results and analysis

In this study total 50 cases were taken. Among them most common age group was 50-60 years comprising 40% of the total. But among female the most common age group was 60-70 years. 72% of patients were from the age range of 50-70years, the most vulnerable age group for stroke. (Figure 1)

52% of patients had hyperlipidemia, 42% had hypertension, 22% smoker, 34% alcoholic and 22% Diabetes. Atrial fibrillation was seen in 18% of cases, more common in female. 10% had CVA in past and 10% of patients were known case of CAD.

Most common symptom at presentation was hemiplegia (92%)(Right sided $>$ left). Facial palsy was the second most common (72%) presenting symptom. Headache, Vomiting and seizure were present in 22%, 4% and 10% of patients respectively. 44% had dysphasia.

Most common artery affected was Middle cerebral artery (44%), followed by lacunar infarction (34%). 12% of patients had bilateral lesion and this affected the prognosis and stroke severity significantly. Size of infarction significantly affected the mortality ($p= 0.000$) and stroke severity ($p= 0.001$).

44% of patients were having very severe stroke with NIHSS score in the range of 20-40. Mean NIHSS in discharged patients was 16.45 whereas mean score of those who died in hospital with stroke was 35.35. Thus NIHSS of the stroke patient at admission significantly affect mortality. ($p=0.00$).

Out of total 50 patients of Acute ischemic stroke (AIS) included in our study, 17 (34%) died. Difference in Mortality between female (5/11) and male (12/34) was not significant. (p= 0.778 by Pearson chi square test).

Age of the patient significantly (p = 0.003) affect the mortality in AIS. The mean age of discharged patients was 55.06 years whereas the mean age of those who died was 65.18 years. Among the various risk factors, important determinants of mortality were: Hypertension (p=0.07), history of prior cerebrovascular accident (p=0.003), unconsciousness at admission (p=0.001) and ischemic changes in their ECG (p=0.000). DM, history of coronary artery disease (CAD), hyperlipidemia, and atrial fibrillation were not contributing to the prognosis/mortality although these are proven risk factor for the occurrence of ischemic stroke.

The mean CRP level of patients included in this study was 21.93 ± 3.60 mg/l. (Table 1) Only 18% of cases were having normal CRP level. 60% of cases were having serum CRP level in the range of 3-30mg/L and 18% were having even the higher (>30mg/l) level maximum being 108mg/l. (Table 2) Age of the patient significantly correlated with the CRP level (Spearman’s correlation co-efficient 0.294 at p-value 0.05). But the sex didn’t affect the rise in CRP (p= 0.490 by student t-test).

Serum CRP level was affecting the stroke severity significantly (with the correlation coefficient >0.643, p-value 0.000). (Figure 2)

The mean CRP level in discharged patients was 10.51 ± 13.21 mg/l whereas it was 44.11 ± 29.7 mg/l in those who died in the hospital. Thus CRP level affected mortality of acute ischemic stroke significantly. (p value < 0.000, independent t-test and pearson chi quare). (Figure 3)

Figure 1 Age and Sex Distribution




Table 2 CRP in Acute Ischemic stroke	
CRP range (mg/L)	Percentage
<3 (normal)	18%
3-10	30%
10-30	30%
>30	22%
Total	100%

Table 1 CRP STATISTICS	
N	50
Mean (mg/L)	21.93
Std. Error of Mean	3.609
Median	13.10
Mode	4
Std. Deviation	25.523
Variance	651.401
Range	106
Minimum	2
Maximum	108

Figure 2: s. CRP vs. Stroke severity (NIHSS)





Figure 3: CRP vs Mortality in acute Ischemic Stroke



V. Discussion

Age is the single most important risk factor for stroke.^[8] The most common age group among patients of 'Acute ischemic stroke (AIS)' in our study was 50-70 years covering 72% of total patients included in the study. (Figure 1) Male to female ratio in our study was ~2.1:1. (Figure 1) US National Stroke Survey observed this as 1.44:1.^[9] 42% of stroke patients in this study had hypertension. 22% patients of acute ischemic stroke were Diabetic in this study. In *Framingham*, persons with glucose intolerance have double the risk of brain infarction of non-diabetic persons.^[10]

Mean CRP in patients of acute ischemic stroke in this study was 21.93mg/l much higher than the normal level of 3mg/l. Median value of CRP was 13.10 and maximum was 108. (Table 1) This signifies the inflammatory reactions as a mechanism of injury during acute ischemic stroke and possible role of anti-inflammatory molecules for the prevention and treatment of infarction. 22% of patients had CRP level above than 30 mg/l on the other hand 18% had normal CRP level.

Winbeck et al observed mean value of CRP in ischemic stroke as 17.5mg/l which corroborates our observations^[12] In the study of Abdulkadir Kocer et al mean CRP in cases of acute ischemic stroke was 31.2 \pm 4.4 whereas in control it was much lower, 3.9 \pm 0.6.^[13] Keith W Muir et al^[14] also observed that mean CRP in cases were much higher than the control.^[14] Rajput et al. had found that among stroke patients from Pakistan, 132 (88%) had elevated CRP (CRP > 10 mg/L).^[5] Moreover, in a study by Di Napoli et al. from Italy, 95 patients (74.2%) with acute ischemic stroke had high CRP levels (> 0.5 mg/dl) at admission.^[16] Muir et al. had detected CRP > 10 mg/L in 96 out of the 228 (42.1%) patients of acute ischemic stroke in UK.^[15]

CRP has evolved from being an association to a risk factor for vascular pathology of heart and brain. Zacho et al., in his population based study, found a high frequency of ischemic heart disease (32%) and ischemic stroke (25%) among patients with high levels of CRP in Denmark.^[18] Ridker et al. from the US, showed high CRP to be a predictor of risk for future myocardial infarction and stroke in healthy men.^[19]

Serum CRP level was affecting the stroke severity significantly (with the correlation coefficient >0.643, p-value 0.000). (Figure 2) Combined p value by ANOVA for this relation is 0.001. In one study, patients with CRP levels \geq 7 mg/L had higher NIHSS scores on admission. Previous studies have shown that patients with increased CRP levels have larger infarctions^[19].

Serum CRP level also correlates with the stroke prognosis. Mean CRP in discharged patient was 10.51mg/l whereas in those who died in hospital it was 44.11mg/l (p value 0.000). (Figure 3) Few studies in China and Nepal had similar findings.^[6] A recent study showed that elevated CRP levels in young patients with ischemic stroke were associated with an increased risk of mortality, even 12 years after the CRP measurements.^[19] Also Idicula et al found a crude association between on admission high CRP and short-term (7 days) functional outcome in patients with acute ischemic stroke.^[7] Song et al in Korea demonstrated that elevated hs-CRP levels on the seventh hospital day, rather than within 24 h after stroke onset, could strongly predict the prognosis of functional disability.^[19] Elevated CRP levels have been associated with poorer outcome in Malaysian patients with acute ischemic stroke.^[14]

VI. Conclusion

Important risk factors for incidence of acute ischemic stroke are advanced age, Hypertension, DM, atrial fibrillation, hyperlipidemia, prior stroke, alcohol and smoking. Determinants of stroke severity are: Age, ischemic changes in ECG, size of infarct and CRP level. Determinants of mortality are: Age, prior CVA, unconsciousness, ischemic changes in ECG, size of infarction, NIHSS score at admission and raised serum level of CRP.

Serum CRP is raised in most of the patients of acute ischemic stroke and its elevation reveals the underlying inflammation causing atherosclerosis as well as CRP as a marker of brain injury. Serum C-reactive protein elevation is associated with more severe stroke and poor prognosis.

References

- [1]. Longo DL, Fauci AS, Kasper DL, Hauser SL, Jameson J, Loscalzo J. eds. Harrison's Principles of Internal Medicine, 19e. New York, NY: McGraw-Hill; 2015.
- [2]. Elias-Smale SE, Kardys I, Oudkerk M, Hofman A, Witteman JC. C-reactive protein is related to extent and progression of coronary and extra-coronary atherosclerosis; results from the Rotterdam study. *Atherosclerosis*. 2007;195:e195–e202.
- [3]. Grau AJ, Bugge F, Ziegler C, Schwarz W, Meuser J, Tasman AJ, et al. Association between acute cerebrovascular ischemia and chronic and recurrent infection. *Stroke*. 1997;28(9):1724–9. [PubMed]
- [4]. Wakugawa Y, Kiyohara Y, Tanizaki Y, Kubo M, Ninomiya T, Hata J, et al. C-reactive protein and risk of first-ever ischemic and hemorrhagic stroke in a general Japanese population: the Hisayama Study. *Stroke*. 2006;37(1):27–32.
- [5]. Rajput MR, Lakhair MA, Shaikh MA, Rind MS, Zafarullah, Bano R. C-Reactive Protein (CRP) and other Risk Factors in Acute Ischemic Stroke Patients. *Journal of Liaquat University of Medical and Health Sciences (JLUMHS)* 2011;10(3):131–3.
- [6]. Huang Y, Jing J, Zhao XQ, Wang CX, Wang YL, Liu GF, et al. High-sensitivity C-reactive protein is a strong risk factor for death after acute ischemic stroke among Chinese. *CNS Neurosci Ther*. 2012;18(3):261–6. [PubMed]
- [7]. Adams HP, Jr, Davis PH, Leira EC, Chang KC, Bendixen BH, Clarke WR, et al. Baseline NIH Stroke Scale score strongly predicts outcome after stroke: A report of the Trial of Org 10172 in Acute Stroke Treatment (TOAST) Neurology. 1999;53:126–131.

- [8]. Brown RD, Whisnant JP, Sicks RD, O'Fallon WM, Wiebers DO. Stroke incidence, prevalence, and survival: secular trends in Rochester, Minnesota, through 1989. *Stroke*. 1996;27:373-380.
- [9]. American Heart Association Heart Disease and Stroke Statistics—2007 Update. 2007. Retrieved July 26, 2007, from www.americanheart.org/downloadable/heart/1166712318459HS_StatsInsideText.pdf.
- [10]. Ralph L. Sacco, Emelia J. Benjamin, Joseph P. Broderick, Mark Dyken, J. Donald Easton, William M. Feinberg, Larry B. Goldstein, Philip B. Gorelick, George Howard, Steven J. Kittner, Teri A. Manolio, Jack P. Whisnant and Philip A. Wolf. *Stroke*. 1997;28:1507-1517, originally published July 1, 1997
- [11]. Banerjee TK, Das SK. Fifty years of stroke researches in India. *Annals of Indian Academy of Neurology*. 2016;19(1):1-8. doi:10.4103/0972-2327.168631.
- [12]. Winbeck K, Poppert H, Etgen T, Conrad B, Sander D. Prognostic relevance of early serial C-reactive protein measurements after first ischemic stroke. *Stroke*. 2002;33(10):2459–64. [PubMed]
- [13]. C-reactive protein is an indicator for fatal outcomes in first-time stroke patients. AbdulkadirKoçer, AbdulkadirKoçer, CüneytCanbulat, ErenGözke, AtillaIhan *Med SciMonit* 2005; 11(11):CR540-544 ID: 430369 Published: 2005-11-01
- [14]. Muir, Keith Wd; Tyrrell, Pippaa; Sattar, Naveedb; Warburton, Elizabethc. Inflammation and ischaemic stroke. *Current Opinion in Neurology*:June 2007 - Volume 20 - Issue 3 - p 334–342 doi: 10.1097/WCO.0b013e32813ba151
- [15]. Muir KW, Weir CJ, Alwan W, Squire IB, Lees KR. C-reactive protein and outcome after ischemic stroke. *Stroke*. 1999;30(5):981–5. [PubMed]
- [16]. Di Napoli M, Papa F, Bocola V. C-reactive protein in ischemic stroke: an independent prognostic factor. *Stroke*. 2001;32(4):917–24. [PubMed]
- [17]. denHertog HM, van Rossum JA, van der Worp HB, van Gemert HM, de Jonge R, Koudstaal PJ, et al. C-reactive protein in the very early phase of acute ischemic stroke: association with poor outcome and death. *J Neurol*. 2009;256(12):2003–8. [PMC free article] [PubMed]
- [18]. Zacho J, Tybjaerg-Hansen A, Jensen JS, Grande P, Sillesen H, Nordestgaard BG. Genetically elevated C-reactive protein and ischemic vascular disease. *N Engl J Med*. 2008;359(18):1897–908. [PubMed]
- [19]. Chaudhuri JR, Mridula KR, Umamahesh M, Swathi A, Balaraju B, Bandaru VCS. High sensitivity C-reactive protein levels in Acute Ischemic Stroke and subtypes: A study from a tertiary care center. *Iranian Journal of Neurology*. 2013;12(3):92-97.