

Alvarado Scoring Systems in Diagnosis of Acute Appendicitis: A Study

*Dr. Manish Verma¹, Dr. Bhavinder K. Arora*²

¹(MS, FAIS, FMAS, FIAGES, Associate Prof., Department of Surgery, Pt B D Sharma PGIMS, Rohtak, India)

²(MS, FAIS, FIAMS, Professor, Department of Surgery, Pt B D Sharma PGIMS, Rohtak, India)

Corresponding Author: * Dr. Manish Verma,

Abstract: The diagnosis of acute appendicitis is mainly based on clinical examination. Various diagnostic scores are available to aid in the diagnosis of acute appendicitis with varying success rates. We assessed Alvarado scoring systems for the diagnosis of acute appendicitis and their negative appendectomy rate in 100 patients of appendicitis. We observed that Alvarado scoring system a good clinical parameter score for diagnosis of acute appendicitis with accuracy rate of 79% in this study. In this study, the sensitivity and specificity value of this score was 82.42% and 44.44% while positive predictive value and negative predictive value of Alvarado score was 93.75% and 20% respectively. The negative appendectomy rate of the Alvarado score was 6.3%.

Keywords: Appendicitis, Appendectomy, Alvarado Score, RIPASA Score.

Date of Submission: 08 -08-2017

Date of acceptance: 23-08-2017

I. Introduction

Acute appendicitis is one of the most common surgical emergencies encountered in the world.[1] The diagnosis of acute appendicitis may be difficult in patients who present with atypical signs and symptoms on physical examination.[2] Alvarado scoring system is one of the most commonly used scoring tool which incorporates symptoms, signs and laboratory investigations to reach the diagnosis.[3] Many studies in the literature are available on diagnostic scores for acute appendicitis in western population but there are only few studies available in Indian population. We carried out a prospective study to assess reliability of Alvarado scoring systems for the diagnosis of acute appendicitis and their negative appendectomy rate.

II. Patients And Methods

The present study was conducted in 100 patients in department of surgery of our institute. All patients presenting with right iliac fossa pain, suspected to be having acute appendicitis were included in this study. Paediatric age group of less than 14yrs were excluded from the study. The data including the patients' demographics central number, age and gender, the presenting symptoms (right illiac fossa pain, nausea and vomiting, anorexia and the duration of symptoms), clinical signs (right illiac fossa tendernes, guarding, rebound tenderness, Rovsing's sign and fever) and laboratory investigations (elevated white cell count and negative urinalysis, shift to left) were recorded. Confirmation of acute appendicitis as the final diagnosis was obtained from histopathological analysis of the resected appendix in the department of pathology. Data regarding patient's admission and discharge dates, date of appendectomy if performed, name and signature of confirming surgeon, post-operative complications and use of radiological investigations were recorded. Data was analysed using SPSS-17 software and appropriate statistical tests (chi Square test, unpaired t test and ANOVA test) were applied to draw conclusion.

III. Observations

In our study, 83% patients were of <40yrs of age and 67% were males. All the patients presented with complaint of pain in Right illiac fossa. Migration of pain to right illiac fossa, anorexia, nausea-vomiting and fever were present in 94%, 88%, 78% and 36% patients respectively. All the patients were having tendernes and rebound tenderness. Guarding and Rovsing's sign were present in 49% and 40% patients respectively. Seventy seven patients had raised leucocyte count with shift to the left seen in only 7 patients. The routine microscopy was normal in 96 patients.

When ALVARADO score was applied to all patients, 80 patients had their score ≥ 7 i.e. 80% patients had acute appendicitis. The scoring system was analyzed with respect to histopathology which is considered the gold standard. A total of 100 patients under study were operated for appendicitis and it was confirmed

histopathologically in 91 patients. Nine patients did not have evidence of appendicitis and were considered to have undergone negative appendectomy as per histopathology. On histopathology, the most common type was acute appendicitis with periappendicitis(49%). Other types were acute diffuse suppurative appendicitis, gangrenous appendicitis, obliterative appendicitis and granulomatous appendicitis. According to Alvarado score, 80 patients were diagnosed to have appendicitis. Out of these 80 patients, only 75 patients had evidence of appendicitis histopathologically. Five patients were falsely diagnosed to have appendicitis by Alvarado scoring system. Out of 20 patients diagnosed by Alvarado as not having appendicitis, 16 patients were missed by this scoring system. However, the findings were not statistically significant (Table no. 1).

Table no. 1: comparison of alvarado score with final histopathological diagnosis
n=100

Alvarado score	Total no of patients	No appendicitis	Appendicitis
Alvarado score <7	20 (100%)	4 (20%)	16 (80%)
Alvarado score ≥7	80 (100%)	5 (6.3%)	75 (93.8%)
Total	100 (100%)	9 (9%)	91 (91%)

Sensitivity of the Alvarado scoring system in the study was 82.42% and specificity came out to be 44.44%. The positive and negative predictive values were 93.75% and 20% respectively. Accuracy of the Alvarado scoring system was 79% in the studied population. The negative appendectomy rate was 6.3% with respect to the histopathology findings (Table no 2).

Table no. 2: alvarado ripasa scoring systems in the diagnosis of acute appendicitis
n=100

Statistical parameters	Alvarado score
Sensitivity	82.42%
Specificity	44.44%
Positive predictive value	93.75%
Negative predictive value	20%
Accuracy	79%
Negative appendectomy rate	6.3%

IV. Discussion

Appendicitis is a condition characterized by inflammation of appendix. Acute appendicitis has a lifetime prevalence rate of approximately 1 in 7.[4] The first description of a case of appendicitis was by Fernel in 1554.[5] McBurney in 1889 contributed his classical sign to the diagnosis of appendicitis.[6] Several other conditions can mimic appendicitis.[7] Only CECT can diagnose the condition with very high sensitivity and specificity but it is not feasible to have this investigation for each and every patient suspected to be having appendicitis, particularly in countries with limited resources.[8,9] Delay in diagnosis can lead to morbidity and even mortality. To prevent this delay, various investigations have been tried but surgeons' good clinical assessment is considered to be the most important requisite in the diagnosis of appendicitis.

Till date we have no laboratory parameters that could indicate or reliably point to presence or absence of acute appendicitis. There has been a need of scoring system with acceptable sensitivity, specificity and negative appendectomy rate. We studied the diagnostic value of Alvarado scoring system for acute appendicitis and analyzed the negative appendectomy rate in 100 patients of appendicitis who underwent appendectomy. In the present study, the mean age of the patients with appendicitis was 28.10±10.887 years. Most of the patients were in the younger age group with age less than 40yrs and male preponderance as evidenced by various studies in the literature.[10-15] The cause of male predominance is still not clear but increased incidence of appendicolith in males is thought to be responsible for it. The most common symptom in patients with appendicitis was pain in the right illiac fossa which is consistent with other studies in the literature.[12,13] Tenderness was observed in all the patients as mentioned in the other studies published in literature.[12,16]

In our study, the most common type of appendicitis on histopathology was acute appendicitis with periappendicitis which was same as reported in other studies in literature. In the present study, negative appendectomy rate was observed in 9% of the patients, which is similar as in the studies done by Erdem et al and Nshuti et al. However, the negative appendectomy rates are variable in different studies in literature.[10,12,13,17-19] In the present study, the sensitivity, specificity, positive predictive value and negative predictive value of Alvarado score was 82.42%, 44.44%, 93.75% and 20% respectively. Other studies in literature show variable results.[10,14,16,20-22] Accuracy in the present study was 79% which was similar to the studies done by Jawaid et al and Erdem et al and lower in the study done by Chong et al. In the present study the negative appendectomy rate of the Alvarado score was 6.3% which was lower than in other studies in the literature available.[16,20,22]

V. Summary And Conclusion

In Summary Alvarado score is a good clinical parameter score for diagnosis of acute appendicitis with accuracy rate of 79% in this study. In this study, the sensitivity and specificity value of this score was 82.42% and 44.44% while positive predictive value and negative predictive value of Alvarado score was 93.75% and 20% respectively. The negative appendectomy rate of the Alvarado score was 6.3%. Our study design should however, be applied to a larger group of patients to further evaluate the validity of our results. This can contribute to the quest for surgical excellence and better patient care for one of the most commonly performed surgical procedure in the world.

References

- [1]. Kumar V, Cotran RS, Robbins SL. Appendix. In: Robbins basic pathology. 5th ed. London: WB Saunders 1992.p.502-2.
- [2]. Gilmore OJ, Browett JP, Griffin PH. Appendicitis and mimicking conditions. A prospective study. *Lancet*; 1975; 2: 421-4.
- [3]. Alvarado A. A practical score for the early diagnosis of acute appendicitis. *Ann Emerg Med*; 1986; 15: 557-64.
- [4]. Stephens PL, Mazzucco JJ. Comparison of ultrasound and the Alvarado score for the diagnosis of acute appendicitis. *Conn Med*; 1999; 63: 137-40.
- [5]. Seal A. Appendicitis: A historical review. *Can J Surg*; 1981; 24: 427-33.
- [6]. McBurney CM. Experience with early operative interference in cases of disease of the vermiform appendix. *N Y Med J*; 1889; 50: 676-84.
- [7]. Graffeo CS, Counselman FL. Appendicitis. *Emerg Med Clin North Am*; 1996; 14: 653-71.
- [8]. Krajewski S, Brown J, Phang PT, Raval M, Brown CJ. Impact of computed tomography of the abdomen on clinical outcomes in patients with right lower quadrant pain: a meta-analysis. *Can J Surg*; 2011; 54(1): 43-53.
- [9]. Ozao-Choy J, Kim U, Menes TS. Incidental findings on computed tomography scans for acute appendicitis: Prevalence, costs and outcome. *Am Surg*; 2011; 77: 1502-9.
- [10]. Dey S, Mohanta Pk, Baruah AK, Kharga B, Bhutia KL, Singh VK. Alvarado scoring in acute appendicitis: A clinico-pathological correlation. *Indian J Surg*; 2010 Aug; 72(4): 290-3.
- [11]. Hale DA, Molloy M, Pearl RH, Schutt DC, Jaques DP. Appendectomy: a contemporary appraisal. *Ann Surg*; 1997 Mar; 225(3): 252-61.
- [12]. Samad A, Kumar B, Khanzada TW, Lohana D. Appropriate cut-off value of Alvarado scores for patients undergoing appendectomy at Isra University Hospital, Hyderabad. *Isra Med J*; 2009 Aug; 1(2): 36-9.
- [13]. Chamisa I. A clinic-pathological review of 324 appendices removed for acute appendicitis in Durban, South Africa; a retrospective analysis. *Ann R Coll Surg Engl*; 2009 Nov; 91(8): 688-92.
- [14]. Alnjadat I, Abdallah B. Alvarado versus RIPASA score in diagnosing acute appendicitis. *RMJ*; 2013; 38(2): 147-51.
- [15]. Nshuti R, Kruger D, Luvhengo TE. Clinical presentation of acute appendicitis in adults at the Chris Hani Baragwanath academic hospital. *Int J Emerg Med*; 2014 Feb 17; 7(1): 12.
- [16]. Erdem H, Centinkunar S, Das K, Reyhan E, Deger C, Aziret M et al. Alvarado, Eskelinen, Ohlmann and Raja Isteri Pengiran Anak Saleha Appendicitis scores for diagnosis of acute appendicitis. *World J Gastroenterol*; 2013 Dec 21; 19(47): 9057-62.
- [17]. Deballon PO, Chalumeau C, Facy O. CT scan and appendicitis: inadequate population. *J Surg Educ*; 2007 Mar-Apr; 64(2): 64.
- [18]. Kanumba ES, Mabula JB, Rambau P and Chalya PL. Modified Alvarado scoring system as a diagnostic tool for acute appendicitis at Bugando Medical Centre, Mwanza, Tanzania. *BMC Surg*; 2011; 11: 4.
- [19]. Emre A, Akbulut S, Bozdog Z, Yilmaz M, Kanlioz M, Emre R et al. Routine histopathological examination of appendectomy specimens: retrospective analysis of 1255 patients. *Int Surg*; 2013 Oct-Dec; 98(4): 354-62.
- [20]. Jawaid A, Asad A, Motiei, Munir A, Bhutto E, Choudary H et al. Clinical scoring system: A valuable tool for decision making in cases of acute appendicitis. *J Pak Med Assoc*; 1999 Oct; 49(10): 254-9.
- [21]. Shrivastava UK, Gupta A, Sharma D. Evaluation of the Alvarado score in the diagnosis of acute appendicitis. *Trop Gastroenterol*; 2004 Oct-Dec; 25(4): 184-6.
- [22]. Chong CF, Thein A, Mackie AJ, Tin AS, Tripathi S, Ahmad MA et al. Comparison of RIPASA and Alvarado scores for the diagnosis of acute appendicitis. *Singapore Med J*; 2011; 52(5): 340-5.

*Dr. Manish Verma. " Alvarado Scoring Systems in Diagnosis of Acute Appendicitis: A Study." IOSR Journal of Dental and Medical Sciences (IOSR-JDMS) 16.8 (2017): 01-03