

Evaluation of Dry Eye: A Hospital Based Study

*Dr Harvinder Nagpal¹, Dr Anil Pathak², Dr Reagan Madan³

¹Assistant professor, Department of ophthalmology, government medical college, Patiala.

²Associate professor, Department of ophthalmology, government medical college, Patiala.

³Senior resident, Department of ophthalmology, government medical college, Patiala.

Corresponding author: Dr Reagan Madan

Abstract: Dry eye is a condition which is normally underdiagnosed. It is a multifactorial disease of tears and ocular surface and varies in severity, duration, with varied etiology. It is not sight threatening but irritating so finding correct factors which causes dry eye needs to be removed as elimination of such factors lead to marked improvement and may cure as it results in symptoms of discomfort, visual disturbance and tear film instability with potential damage to ocular surface. So doing tests to properly assess and grade and treat accordingly is the need of the hour because if not properly diagnosed and treated it can cause sight threatening complications.

Background and objectives: A prospective study was conducted to assess the dry eye in a tertiary care hospital of north india. Along with assessment of dry eye with OSDI questionnaire and various tests were done to grade dry eye.

Methods: 63 cases were chosen from the outpatient department of and assessment of dry eye wasmade by tests like Schirmer's test, Tear breakup time, Rose Bengal dye test. Ocular surface disease index (OSDI) questionnaire was given to patients to assess the grading of dry eye.

Results: Majority 31.7% patients were in the group 41-50 years. In the study OSDI questionnaire had a good reliability and consistency ($p < 0.001$) highly significant. Pearson correlation with r value among various test like Rose Bengal test, Schirmer's test and TBUT showed a good correlation.

Conclusion : Dry eye is a chronic disease and increase in incidence of dry eye increases with age. Various risk factors should be assessed to find the cause of dry eye. Subjective test like Odi questionnaire with objective tests like rose Bengal, Schirmer's and Tear breakup time should be done routinely to assess the grading of dry eye so that long term complications associated can be prevented.

Keywords: OSDI; Schirmer's ; Rose Bengal; Tear break up time

Date of Submission: 20 -09-2017

Date of acceptance: 30-09-2017

I. Introduction

Dry Eye is a common condition that is often under diagnosed. Normal vision requires moist healthy ocular surface. A sufficient quality of tears, normal composition of tears film, lid closure to maintain healthy ocular surface.⁽¹⁾Dry eye is a disorder characterised by either quantitative decrease or qualitative change in precorneal film resulting in spectrum of pathological changes that may adversely affect the ocular surface resulting in ocular surface disorders often leading to conjunctival squamous metaplasia and punctate epithelial erosion of cornea.^(2,3)Dry eye results in discomfort and visual disturbance and tear film Instability with potential damage to ocular epithelial surface and accompanied by increase in tear osmolarity and inflammation. Dry eye syndrome involves multiple risk factors that when disregarded can result in treatment failure and frustration both for the patients and the physician. Dry eye may lead to increase risk of infections, medications toxicity, contact lens intolerance, progressive ocular surface disease, scarring, cornea morbidity namely keratinisation, corneal thinning, vascularisation, microbial and sterile corneal ulcer leading to perforation and severe visual loss. Hence correct diagnosis and appropriate management of dry eye is essential.⁽⁴⁾

Dry eye etiology is not a single disease entity. It is a disorder of lacrimal function unit according to some. In recent population based surveys it indicates that dry eye affects millions of people in worldwide as many as 20-25% patients reported to out patient department complaints of dry eye symptoms making it very common presentation. It poses an huge economic burden to patients if not evaluated for risk factors, etiopathogenesis that may result in delay in treatment. It affects the quality of life of patients with regards to daily visual acuity. It results in social stigma as a patient may have chronic red eye. Patient may undergo into depression which requires additional psychotherapy. The purpose of approach to is to assess dry eye with OSDI questionnaire along with battery of tests so as to improve patients comfort and prevent structural damage to ocular component. Due to lack of uniformity in definition and inability of any single diagnostic test or set of diagnostic test to confirm or rule out the condition.⁽⁵⁾There has been a shift towards symptom based assessment as a key component in clinical diagnosis with grading of severity of dry eye.

Use of symptom based validated questionnaire might be beneficial as it allows grading of symptoms and is repeatable for comparative purpose before, during and after treatment. Recent advances in treatment suggests the use of lubricants, anti-inflammatory drugs, plugs to augment the tear film.⁽⁶⁾

II. Materials And Methods

The material for the study was collected from the patients presenting themselves directly to Department of Ophthalmology in tertiary hospital of north india.

Inclusion criteria:

Both male and female patients equal to and above 20yrs presenting with following symptoms Burning sensation, sandy gritty feeling, Foreign body sensation, Photophobia, Heavy lids. The above symptoms increase in conditions of low humidity and wind.

Exclusion criteria:

1. Patients less than 20yrs of age, Patients with H/O increased mucoid discharge and watery secretion suggestive of vernal keratoconjunctivitis, Patients with H/O Alkali burns, Patients with H/O Trachoma, Patients with H/O Acute ocular infections, Patients with H/O Ocular surgery within last 6 months, Patients with H/O Impaired eyelid function like in Bells palsy, nocturnal lagophthalmos, ectropion and Contact lens users

An OSDI-Allergan ocular surface disease index questionnaire is to be administered to all participants to assess the symptoms of dry eye and correlate them with OSDI (provided by Allergan, Inc. Irvine, CA) was used to quantify the specific impact of dry eye.⁽⁷⁾ This disease-specific questionnaire includes three subscales: ocular discomfort (OSDI-symptoms), which includes symptoms such as gritty or painful eyes; functioning (OSDI-function), which measures limitation in performance of common activities such as reading and working on a computer; and environmental triggers (OSDI-triggers), which measures the impact of environmental triggers, such as wind or drafts, on dry eye symptoms. The questions are asked with reference to a one-week recall period. Possible responses refer to the frequency of the disturbance: none of the time, some of the time, half of the time, most of the time, or all of the time. Responses to the OSDI were scored using the methods described by the authors. Subscale scores were computed for OSDI-symptoms, OSDI-function, and OSDI-triggers, as well as an overall averaged score. OSDI subscale scores can range from 0 to 100, with higher scores indicating more problems or symptoms. However, we subtracted the OSDI overall and subscale scores from 100, so that lower scores would indicate more problems or symptoms.⁽⁸⁾

- A complete slit-lamp examination of lid margin, tear meniscus, conjunctiva, cornea and tear film is done. relevant examination of other important ocular structure is done


Fallowing tests to diagnose dry eye syndrome

1. Tear break up time (TBUT)
2. Rose Bengal staining
3. Schirmer's test

Diagnosis and confirmation of dry eye was done by series of test, which in standard order of eye examination are as follows: Tear film break up time (TBUT), slit lamp examination of the anterior segment, assessment of the meibomian glands and schirmer-1 test and lastly the Rose Bengal staining. TBUT was done first because manipulation of the eyelid may affect the result. The test was repeated 3 times in each eye and the average time was recorded. The test is considered positive if the average TBUT is <10 seconds in 1 or both the eyes. To determine the condition of the meibomian glands, the eyelid margins in both the lower and upper lids was examined in the slit lamp. Digital pressure was applied on the tarsi to assess the degree of obstruction. The presence of lid margin telangiectasia, collarette and meibomian gland plugging was recorded and graded. After a minimum gap of 30 minutes, Schirmer-1 test was performed.


A precalibrated dry filter paper strip (whatman filter paper no -41) was placed in each lower fornix at the junction of outer and middle thirds without touching the cornea and left for 5 minutes and patient was asked to close his or her eyes. After 5 minutes the strips was removed and the amount of wetting in mm was recorded. The result was considered positive if the amount of wetting of the paper is <5mm. Rose Bengal staining was done again after 30 minutes, taking care to avoid touching the ocular surface. A van Bijsterveld's score of 4 or more was considered positive for dry eye diseases. All dry eye cases were given appropriate treatment and the follow up was done upto 3 months.

Table 1 : Age distribution



It is observed from the present study that majority 20 patients (31.7%) were in the age group 41-50 years. The mean age in the present study group was (40.8) years with standard deviation of (11.5) years.

Table 2: Age Distribution Of Patients Studied In Association With Prevalence Of Dry Eye



In the present study prevalence was highest in the 41-50 age group of which 2(22.2%) were normal, 2(12.5%) had mild dry eye,10(43.5%) had moderate dry eye,6(40%) had severe dry eye.

Table 3: TBUT Test Of Patients Studied In Association With Prevalence Of Dry Eye

TBUT Test	Dry Eye				Total	P value
	Normal	Mild	Moderate	Severe		
Right Eye	11.00±2.45	9.50±2.00	6.39±2.35	3.20±1.15	7.08±3.40	<0.001**
Left Eye	12.00±2.96	10.38±2.42	7.09±2.48	3.67±1.05	7.81±3.68	<0.001**

In the present study TBUT of patient studied in association of dry eye showed right eye mean of 9.5 ± 2standard deviation in mild dry eye group, mean of 6.39±2.35 standard deviation in moderate dry eye group, mean of 3.20 ±1.15 standard deviation in the severe dry eye group showing a significant p value of less than 0.001. Whereas in the left eye mean of 10.38±2.42 standard deviation in the mild dry eye group, mean of 7.09±2.48 standard deviation in the moderate dry eye group, mean of 3.67± 1.05 standard deviation of in severe dry eye group showing a p value of less than 0.001.

Table 4: Schirmer’s Test Of Patients Studied In Association With Prevalence Of Dry Eye

Schirmer’s test	Dry Eye				Total	P value
	Normal	Mild	Moderate	Severe		
Right Eye	12.33±3.00	10.19±2.81	8.26±3.40	4.47±0.83	8.43±3.76	<0.001**
Left Eye	13.33±4.36	10.88±3.16	8.91±3.59	4.87±1.30	9.08±4.19	<0.001**


In the present study schirmer’s test of patient studied with the prevalence of in right eye mean of 10.19 +2.81 of standard deviation in the mild dry eye group, mean 8.26 with ±3.40 of standard deviation in the moderate dry eye group, mean of 4.47 with ±0.83 standard deviation in the severe dry eye group. For left eye mean of 10.88 with ±3.16 of standard deviation in the mild dry eye group, mean of 8.91 with ±3.59 of standard deviation in the moderate dry eye group, mean of 4.47 ± 1.30 of standard deviation in the severe dry eye group with p value for both eyes being highly significant p<0.001.

Table 5: Rose Bengal Test (Bjsterveld’s Score) Of Patients Studied In Association With Prevalence Of Dry Eye

ROSE BENGAL TEST	Dry Eye				Total	P value
	Normal	Mild	Moderate	Severe		
Right Eye	2.89±1.05	4.19±1.52	5.22±1.31	7.47±0.83	5.15±1.94	<0.001**
Left Eye	2.89±1.16	4.12±1.59	5.22±1.53	7.67±0.82	5.19±2.07	<0.001**

In the present study Rose Bengal test of patient studied with the prevalence of in right eye mean of 4.19±1.52 standard deviation in the mild dry eye group, mean of 5.22 with ±1.31 standard deviation in the moderate dry eye group, mean of 7.47 with ±0.83 standard deviation in the severe dry eye group. For left eye mean of 4.12 with ±1.59 standard deviation in the mild dry eye group, mean of 5.22 with ±1.53 of standard deviation in the moderate dry eye group, mean of 7.67 with ± 0.82 standard deviation in the severe dry eye group. with p value for both eyes being highly significant p<0.001.(ANNOVA TEST).

Table 6: OSDI Score Before And After Treatment



In the present study of assessing OSDI on scale of 0-100 shows scale of <40 before treatment group as 27(42.9%), after treatment 47(74.6%), having a percentage change of 31.7%. In the scale of 40-80, 31(49.2%) before treatment, 15(23.8%) after treatment, having a percentage change of -25.4%. In the scale of >80,5 (7.9%) before treatment group,1(1.6%) after treatment group showing a percentage change of -6.3%.

Table 7: OSDI score and Scale Before Treatment of patients studied in association with prevalence of dry eye

Before Treatment	Dry Eye				Total	P value
	Normal	Mild	Moderate	Severe		
OSDI Score	11.32±4.49	23.79±5.37	42.15±9.63	75.81±11.14	41.10±23.84	<0.001**
OSDI Scale	12.97±1.55	25.04±7.65	42.81±10.46	75.85±10.63	41.90±23.51	<0.001**


the present study of patients before treatment in association with the prevalence of dry eye shows OSDI score of 41.10 mean with ±23.84 standard deviation, on OSDI scale 41.90 with ±23.51 standard deviation both showing p value less than (p<0.001) highly significant.

Table 8: OSDI score and Scale after Treatment(end of 3 months follow up) of patients studied in association with prevalence of dry eye

After Treatment	Dry Eye			Total	P value	
	Normal	Mild	Moderate			
OSDI Score	10.86±4.57	14.61±4.07	28.24±7.07	56.71±13.49	29.08±18.78	<0.001**
OSDI Scale	12.97±1.55	13.73±5.66	28.17±5.63	56.81±13.31	29.15±18.55	<0.001**

In the present study prevalence of dry eye patients after treatment was associated with OSDI score mean of 29.08 with ±18.78 standard deviation and scale showed mean of 29 with ±18.55 standard deviation both showing p value less than (p<0.0001) highly significant.

Table 9: Prevalence Of Dry Eye In Patients Studied



In the present study prevalence of dry eye was noted as follows mild dry eye 16(25.4%), moderate dry eye 23(36.5%), severe dry eye 15(23.8%).

Table 10: Pearson Correlation

Pair	Pearson Correlation	
	R Value	P Value
Osdi Score Vs Tbut	-0.801	<0.001**
Osdi Score Vs Schirmers Test	-0.695	<0.001**
Osdi Score Vs Rose Bengal Test	0.772	<0.001**
Tbut Vs Schirmers Test	0.833	<0.001**
Tbut Vs Rose Bengal Test	-0.817	<0.001**
Schirmers Test Vs Rose Bengal Test	-0.757	<0.001**

In the present study correlation among the various test like TBUT, Schirmer's, Rose Bengal test, OSDI showed good correlation with significant p value among all p<0.001 highly significant. In our study comparison of variable according to the prevalence of dry eye with regards to the OSDI, TBUT, Schirmer's, rose Bengal test reveals strongly significant p value. OSDI mean 41.10±23.84 standard deviation, TBUT mean 7.44±3.51 standard deviation, schirmer's test mean 8.75±3.92 standard deviation, Rose Bengal mean 5.17±1.98 standard deviation overall validated the tests done.

III. Discussion

The present study revealed that there is appreciable variation among diagnostic test results among different diseases and the best test combination to detect dry eye is OSDI/TBUT/Schirmer/Rose Bengal test. This result emphasises the importance of most commonly used tests to detect dry eye in clinical practice and also their variability. Meaningful diagnostic testing in dry eye disease across a broad range of different aetiologies and presentations is still a challenge⁽⁹⁾. Owing to great variability in dry eye disease severity, it is unlikely that a single test result has adequate sensitivity to serve for dry eye given its multifactorial nature and numerous manifestations. So that's why research on potential dry eye diagnostic tools and therapeutic agents has increased exponentially.

The correlation between subjective and objective findings was good but not in all cases and could be caused by multifactorial nature of dry eye syndrome.

In our study 50 (79.4%) of patients have some symptoms such as foreign body sensation, burning, grittiness related to dry eye. there was good association between subjective symptoms of dry eye and its validation with objective test like, Tbut, Schirmer's, Rose Bengal test⁽¹⁰⁾. Objective studies of dry eye commonly involved TBUT, Schirmer's, Rose Bengal test (bjisterveld's score) and when used in our study showed significant p value of less than 0.001.

Also in the present study the OSDI demonstrated consistency and good test reliability. The OSDI also demonstrated excellent validity effectively discriminating between normal, mild, moderate and severe dry eye diseases as defined by both the physician assessment of severity and composite disease severity score.⁽¹¹⁾ Pearson correlation with r value shows perfect correlation between test and significant p value(0.001)OSDI score/scale showed strongly significant value with dry eye before and after treatment (p less than 0.001). OsdI was very well correlated with the clinical tests done suggesting that it is a very good tool if answered properly by the patients and can give the status of dry eye before moving to any tests.

IV. Conclusion

Consensus regarding protocol in dealing with dry eye clinical diagnosis and standardisation of objective test is lacking. Dry eye is a complex multifactorial disease that involves the eyelids, tear film, ocular surface, immune and autonomic nervous system. The condition may be asymptomatic but often time causes significant patient discomfort and visual degradation. More studies into the association of risk factors in dry eye causation needs to be done. Finding an optimal treatment regimen can often be challenging and is a condition that requires patience from both the physician and patient. While mild case require annual visit, most patients suffering from moderate to severe dry eye require frequent monitoring for potential complications of dry eye and treatment efficacy. However the successful treatment of dry eye can lead to significant patient satisfaction and improved quality of life.

It is recommended that establishment of dry eye clinics at rural health centres should be done. Dry eye disease evaluation should be part of eye examination routinely. OSDI questionnaire should be part of working protocol in evaluating dry eye TBUT, Schirmer's, Rose Bengal test which are objective test when done correctly correlates well with OSDI subjective test.

References

- [1]. Begley C G et al. The Relationship between habitual patient reports symptoms and clinical signs among patients with dry eye of varying severity. Invest Ophthalmol Vi Sci 2003;44:4753-61.
- [2]. Elmer Y et al. Dry Eye. Yanoff and Duker Ophthalmology 3rd edition. Mosby Elsevier Publishers 2009;3:324-9.
- [3]. International Dry Eye Workshop Subcommittee. The Definition and Classification of Dry Eye Disease Workshop DEWS. Ocul Surf 2007;5:75-97.
- [4]. Plugfelder SC, Stephen C, Solomon, Stern, Michael. The diagnosis and management of dry eye 25 years review. Cornea 2000; 19:644-99.
- [5]. Afonso AA. Correlation of tear fluorescein clearance and Schirmer's test score with ocular irritations symptoms. Ophthalmology 1999; 106:803-10.
- [6]. Schaumberg DA, Julie E. Buring, David A.Sullivan, Reza Dana. Hormone replacement therapy and dry eye syndrome. JAMA 2001; 286:2114-9.
- [7]. Keith MS, Sudharshan L, Snedecor SJ. Associations between signs and symptoms of dry eye disease: a systematic review. Clin Ophthalmol 2015; 9:1719-30. .
- [8]. Mizuno Y et al. Association between clinical diagnostic tests and health-related quality of life surveys in patients with dry eye syndrome. Jpn J Ophthalmol. 2010; 54(4):259-65.
- [9]. Sullivan BD et al. Correlations between commonly used objective signs and symptoms for the diagnosis of dry eye disease: clinical implications. Acta Ophthalmol.
- [10]. Golding TR et al. Diagnostics Accuracy and intercorrelation of clinical tests for dry eye. Invest Ophthalmol Vis Sci 1993; 34:823.
- [11]. Grubbs JR, Tolleson-Rinehart S, Huynh K, Davis Rm. A review of quality of life measures in dry eye questionnaires. Cornea. 2014 Feb; 33(2):215-8.

*Dr Harvinder Nagpal. "Evaluation of Dry Eye: A Hospital Based Study." IOSR Journal of Dental and Medical Sciences (IOSR-JDMS) 16.9 (2017): 44-49