

## Gingival Recession in a Rural Community in Nigeria: A Pilot Study

Dr. OSADOLOR O.O<sup>1</sup>, Dr. Akaji E.A<sup>2</sup>, Dr. Onyejaka N.K<sup>3</sup>, Dr. Osadolor A.J<sup>4</sup>

<sup>1</sup>Department of Child Dental Health, University of Nigeria Teaching Hospital, Ituku- ozalla, Enugu State, Nigeria.

<sup>2</sup>Department of Preventive Dentistry, University of Nigeria, Enugu State.

<sup>3</sup>Department of Child Dental Health, University of Nigeria, Enugu State.

<sup>4</sup>Central hospital, Benin-city. Edo State.

Corresponding author: Dr. OSADOLOR O.O

---

**Background:** Gingival recession could affect an individual's quality of life with nutritional, functional and psychosocial consequences when associated with clinical problems such as root surface hypersensitivity, root caries, cervical root abrasions, erosions, calculus/plaque retention and chronic periodontitis.

**Objective:** To determine the prevalence of gingival recession in a rural community in South-east Nigeria.

**Methods:** A cross sectional descriptive study of 54 participants was carried out at Ozalla Model Primary Health Centre. They were interviewed for socio-demographic characteristics using interviewer administered questionnaire and intra-oral examination using disposable instruments for each participant was conducted to determine the presence or absence of gingival recession. Ethical clearance for this study was sought and obtained. After data collection, oral health education and tooth brushing demonstrations were done. Data was analysed using Statistical Package for Social Sciences (SPSS) Version 20. P values < 0.05 were accepted as being statistically significant.

**Results:** A total of 54 participants were seen and examined, 27(50 %) were males and 27(50%) were females with male to female ratio of 1:1. The age range of participants examined was 17- 84 years and the mean age was 44.5± 17.4 years. The prevalence of gingival recession was 22.2 %. Majority (85.2%) of the participants uses toothbrush with toothpaste only as an oral hygiene device, 81.5% of the participants brush their teeth once daily and 53.7% had not visited a dental clinic. P values < 0.05 were accepted as being statistically significant. P = 0.001.

**Conclusions:** The prevalence of gingival recession was 22.2 %. Gingival recession was more common in mandibular teeth and both sides of the dental arch.

**Key words:** Gingival recession, rural, community

---

Date of Submission: 13-05-2020

Date of Acceptance: 26-05-2020

---

### I. Introduction

Gingival recession could affect an individual's quality of life with nutritional, functional and psychosocial consequences when associated with clinical problems such as root surface hypersensitivity, root caries, cervical root abrasions, erosions, calculus/plaque retention and chronic periodontitis. For a patient, gingival recession usually creates an aesthetic problem, especially when it affects the anterior teeth [1], and anxiety about tooth loss due to progressing loss of tooth support. Gingival recession may be localized to one tooth, or a group of teeth, or may be generalized throughout the oral cavity. It may be associated with apical shift of marginal gingiva on one or more surfaces resulting in clinical attachment loss and root exposure.[2]

The purpose of using tooth cleaning aids like chewing stick, and toothbrush with toothpaste is to attain good oral hygiene compatible with good oral health. Unfortunately, the practice when not performed appropriately, may result into unwanted ill effects such as recession of the gingival tissues, trauma to soft tissues and abrasion of dental hard tissues [3].

The causes of gingival recession is multifactorial and several predisposing factors may play a role in gingival recession development, such as vigorous teeth brushing, destructive periodontal disease, tooth malposition, frenum pull ,chewing stick trauma, occlusal trauma and iatrogenic factors like inappropriate fixed prostheses, poorly designed partial dentures, operative procedures, and some orthodontic treatment.[4-6]

Mohamed et al., examined the relationship between chewing sticks (Miswak) and gingival recession and he reported that The Miswak users had significantly more sites with gingival recession than did the toothbrush users [7]. He concluded that the Miswak should be considered as a possible factor in gingival recession.

Previous epidemiological studies in Tanzania have shown that Tooth cleaning devices commonly used include plastic toothbrushes (52-68%), chewing sticks (26-32%), both plastic toothbrushes as well as chewing sticks (17.0%), and dental floss (1%) [8-9]. Most of the rural population that constitutes about 85% of the general population in Tanzania uses chewing sticks rather than the toothbrush (8-9).

Some studies reported that the maxillary canines and premolars [10] were the teeth most frequently affected by gingival recession, others reported that the mandibular lateral incisors and premolars, the maxillary and mandibular first molars are commonly affected [11]. Sarpangala et al., in 2015 [12] reported that canines of both the upper right and left were the most frequent regions affected by gingival recession. It has been discovered that the distribution pattern of gingival recessions is related to different etiologic factors [11] Gingival recessions on the mandibular incisors were tied to poor oral hygiene [13] whereas those on the premolars were linked to traumatic tooth brushing [14]

A high prevalence of gingival recession has been reported in America (63%-89%) [15-17], Europe (25%-84%) [18-23] and Australia (71%) [24] but a lower prevalence has been found in Africa (28%) [14, 25,] and Asia (15%) [26]. Humagain and Kafle reported [2] a prevalence of 65.44% from a study conducted in Nepal, India. Mumghamba *et al.*, from reports in East-Africa, conducted a study in Tanzania and reported a prevalence of 33.6% [3]. Arowojolu reported [25] a prevalence of 27.7% from a study done in Ibadan, Nigeria.

There are various studies and reports of the prevalence of gingival recession in other parts of the country and the World. The aim of this study is to determine the prevalence of gingival recession in a rural community in South-east Nigeria in order to contribute to the existing data on gingival recession in Nigeria and the West African sub-region. It will also compare findings with published reports from Nigeria and other countries of the World.

## **II. Materials And Method**

The study was conducted among participants who presented at Ozalla model primary health centre at the time of the study. Ozalla town is one of the thirty three (33) towns that make up Nkanu -West local Government Area of Enugu State [27]. The area is largely rural and its inhabitants are primarily farmers, small-scale business people and traders. Nkanu-West Local Government area of Enugu State is one of the seventeen (17) local government areas of Enugu State and has its headquarters at Agbani [27]. Ethical clearance for this study was sought and obtained. Permission was sought from the head of the community and health centre, while individual verbal consent was obtained from the respondents before giving out the pre-tested questionnaire which was interviewer-administered and an intra oral examination using disposable instruments for each person examined was conducted for the presence/absence of gingival recession. After data collection, oral health education and tooth brushing demonstrations were done. Data were analysed using a computer software programme, Statistical Package for Social Sciences (SPSS) Version 20. P values < 0.05 were accepted as being statistically significant.

## **III. Results**

A total of 54 participants were seen and examined, 27(50 %) were males and 27(50%) were females with male to female ratio of 1:1. The age range of the participants examined was 17years to 84 years with a mean age of  $44.5 \pm 17.4$  (**Table 1**) 12 (22.2%) of the participants presented with gingival recession, 42 (77.8%) of the participants did not present with gingival recession. The participants who presented with gingival recession were 9 males and 3 females with a ratio of 3:1[M: F] (**Table 3**). The predominant occupation was farming and small-scale business/ trading. 85.2% of the participants uses toothbrush with toothpaste only as oral hygiene device.(**Table 2**), 81.5% of the participants brush their teeth once daily and 53.7% had not visited a dental clinic. Mandibular teeth were more commonly affected with gingival recession.(**Figure 1**).

**TABLE 1: SOCIO-DEMOGRAPHIC CHARACTERISTICS OF RESPONDENTS**

Variable	Frequency	Percent
Gender		
Male	27	50
Female	27	50
Age group		
10-19	2	3.7
20-29	12	22.2
30-39	10	18.5
40-49	8	14.8
50-59	6	11.1
60-69	10	18.5
70-79	5	9.3
80-89	1	1.9
Level of education		
Primary	15	27.8
Secondary	24	44.4
Tertiary	15	27.8
Total	54	100

**Table 2: Oral hygiene device used by the participants**

Oral hygiene device used	Frequency	Percent
Toothbrush	46	85.2
Chewing stick	4	7.4
Toothbrush& chewing stick	4	7.4
Total	54	100

**Table 3 Age and gender relationship of participants with gingival recession**

Variable	Frequency	Percent
Gender		
Male	9	75
Female	3	25
P = 0.051		
Age group		
20-40	3	25.0
50-60	3	25.0
Over 60	6	50.0
P- value = 0.001		
Total	12	100




Figure 1: Teeth with gingival recession




Figure 2: Side of the dental arch with presence of gingival recession

#### IV. Discussion

Gingival recession has been defined as a displacement of gingival margin apically from the cemento-enamel junction (CEJ), leading to root-surface exposure [1]. It is commonly observed in adult subjects and can be localized or generalized [2]. The prevalence of gingival recession was 22.2 % of the studied population, which was slightly less than previous studies ranging from 22.5% to 28 % [25,28] . This study demonstrated higher prevalence of gingival recession in mandibular teeth as compared to maxillary teeth, which is in agreement with the findings of previous studies.[8,11, 14,29] . Lower occurrence of gingival recessions in maxillary teeth is probably related to the characteristics of keratinized mucosa, which is wider and thicker in maxilla than in the mandible.[22]

The occurrence of gingival recession correlated with age, which is supported by reports of previous studies [14,17,25 30]. This study confirms that the prevalence of gingival recession increase with age; which is

consistent with most of the epidemiological studies on several age groups.[2,31-32] . The relationship between increased prevalence of gingival recession and age could be due to the cumulative effect of age, periodontal disease and longer period of exposure to the agents that cause gingival recession [2]. Majority (85.2%) of the participants used a plastic toothbrush, few used chewing sticks (7.4%) and a combination of chewing sticks and plastic toothbrushes (7.4%). This study showed greater gingival recession in males than in females which was similar with previous reports[1,33]

## V. Conclusion

The prevalence of gingival recession was 22.2 %. It was more common in mandibular teeth and both sides of the dental arch. The occurrence of gingival recession increases with age and was more common in males than females.

## Financial support and sponsorship

None

## Conflicts of interest

There are no conflicts of interest

## References

- [1]. Khosya B, Devaraj CG. Etiology and severity of different grades of gingival recession in adult population .National Journal of Medical Research,2014, Volume 4,Issue 3.
- [2]. Humagain M, Kafle D. The Evaluation of Prevalence, Extension and Severity of Gingival Recession among Rural Nepalese Adults, Orthodontic Journal of Nepal, 2013,Vol. 3, No. 1.
- [3]. Mumghamba EGS, Honkala S, Honkala E , Manji KP. Gingival Recession, Oral Hygiene and Associated Factors Among Tanzanian Women. East African Medical Journal ,2009,Vol. 86. No. 3.
- [4]. Tugnait A, Clerehugh V. Gingival recession: its significance and management.Journal of Dentistry. 2001; 29: 381-394.
- [5]. Kundapur PP, Bhat KM, Bhat GS. Association of Trauma from Occlusion with Localized Gingival Recession in Mandibular Anterior Teeth. DRJ 2009; 6(2): 71-4.
- [6]. Greenwell H, Fiorellini J, Giannobile W, Offenbacher S, Salkin L, Townsend C, et al. Oral reconstructive and corrective considerations in periodontal therapy. J Periodontol 2005;76(9): 1588-1600.
- [7]. Mohamed AE, Abdullah R, Hassan AS. The relationship between chewing sticks (Miswak) and periodontal health. III relationship to gingival recession. Quintessence International 1991 ; 21 (1) : 61-64.
- [8]. Mumghamba EGS, Fabian FM. Periodontal health status and oral hygiene practices among middle-aged adults in Mtwara rural, Tanzania. Afr.J. Oral. Health Sci. 2003; 4: 192-197.
- [9]. Sarita PTN, Tuominen R. Tooth cleaning methods and their effectiveness among adults in rural Tanzania. Proc. Finn. Dent. Soc. 1992; 88: 139-145.
- [10]. Watson PJC. Gingival Recession. Journal of Dentistry.1984; 12(1):29-35.
- [11]. Anarthe R, Mani A, Marawar PP . Study to Evaluate Prevalence, Severity and Extension of Gingival Recession in the Adult Population of Ahmednagar District of Maharashtra State in India. IOSR Journal of Dental and Medical Sciences (IOSR-JDMS).2013; 6 (1): 32-37.
- [12]. Sarpangala M, Suryanarayan MA, Shashikanth H, Shanker KR, Mohamed M, Devasya A. A study on etiology and occurrence of gingival recession - An epidemiological study. Indian Soc Periodontol. 2015;19(6):671-675.
- [13]. Addy M, Mostafa P, Newcombe RG .Dentine hypersensitivity: the distribution of recession, sensitivity and plaque. Journal of Dentistry.1987; 15(6): 242-248.
- [14]. Van Palenstein Helderman WH, Lembariti BS,van der Weijden GA, van't Hof MA. Gingival recession and its association with calculus in subjects deprived of prophylactic dental care. Journal of Clinical Periodontology.1998; 25(2): 106-111.
- [15]. Khocht A, Simon G, Person P, Denepitiya JL. Gingival recession in relation to history of hard toothbrush use. Journal of Periodontology. 1993; 64: 900-905.
- [16]. Marini MG, Gregghi SL, Passanezi E, Sant'ana AC. Gingival recession: prevalence, extension and severity in adults. Journal of Applied Oral Science. 2004; 12: 250-255.
- [17]. Susin C, Haas AN, Oppermann RV, Haugejorden O, Albandar JM. Gingival recession: epidemiology and risk indicators in a representative urban Brazilian population. Journal of Periodontology. 2004; 75: 1377-1386.
- [18]. Checchi L, Daprile G, Gatto MR, Pelliccioni GA. Gingival recession and toothbrushing in an Italian School of Dentistry: a pilot study. Journal of Clinical Periodontology.1999; 26: 276- 280.
- [19]. Kozłowska M, Wawrzyn-Sobczak K, Karczewski JK, Stokowska W. The oral cavity hygiene as the basic element of the gingival recession prophylaxis. Roczniki Akademii Medycznej w Białymstoku. 2005; 50 Suppl 1: 234-237.
- [20]. Sarfati A, Bourgeois D, Katsahian S, Mora F, Bouchard P. Risk assessment for buccal gingival recession defects in an adult population. Journal of Periodontology.2010; 81: 1419- 1425.
- [21]. Serino G, Wennstrom JL, Lindhe J, Eneroth L. The prevalence and distribution of gingival recession in subjects with a high standard of oral hygiene. Journal of Clinical Periodontology. 1994; 21: 57-63.
- [22]. Tenenbaum H. A clinical study comparing the width of attached gingiva and the prevalence of gingival recessions. Journal of Clinical Periodontology. 1982; 9: 86-92.
- [23]. Toker H, Ozdemir H. Gingival recession: epidemiology and risk indicators in a university dental hospital in Turkey. International Journal of Dental Hygiene. 2009; 7: 115-120.
- [24]. Thomson WM, Broadbent JM, Poulton R, Beck JD. Changes in periodontal disease experience from 26 to 32 years of age in a birth cohort. Journal of Periodontology. 2006; 77: 947-954.
- [25]. Arowojolu MO. Gingival recession at the University College Hospital, Ibadan: prevalence and effect of some aetiological factors. African Journal of Medicine and Medical Sciences. 2000; 29: 259-263.
- [26]. Slutzkey S, Levin L. Gingival recession in young adults: occurrence, severity, and relationship to past orthodontic treatment and oral piercing. American Journal of Orthodontic and Dentofacial Orthopedics. 2008; 134: 652-656.

- [27]. Adekola G, Nwoye CE. Traditions and Customs in Community Development: The Case of Nkanu West and Nkanu East Local Government Areas of Enugu State, Nigeria. *Journal of Education and Practice*. 2016: Vol.7, No.18.
- [28]. Muller HP, Stadermann S, Heineeke A. gingival recession in smokers and non-smokers with minimal periodontal disease. *J Clin Periodontol* 2002 Feb; 29(2): 129-36.
- [29]. Vehkalahti, M. Occurrence of gingival recession in adults. *J. Periodontol.* 1989; 60: 599-603
- [30]. Mumghamba, E.G.S., Markkanen, H.A. and Honkala, E. Risk factors for periodontal diseases in Ilala, Tanzania. *J. Clin. Periodontol.* 1995; 5: 347-354.
- [31]. Anerud A, Boysen H. The natural history of periodontal disease in man: prevalence, severity and extent of gingival recession. *J Periodontol* 1992 Jun; 63: 489- 495
- [32]. Guimaraes MG, Aguiar EG. Prevalence and type of gingival recession in adults in the city of Divinopolis, MG, Brazil. *Braz J Oral Sci* 2012; 11(3): 357-361.
- [33]. Dodwad V. Aetiology and severity of gingival recession among young individuals in Belgaum district in India. *Annal Dent Univ Malaya* 2001; 8: 1-6.

Dr. OSADOLOR O.O, et. al. "Gingival Recession in a Rural Community in Nigeria: A Pilot Study." *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)*, 19(5), 2020, pp. 34-39.