

Chemical Composition And Larvicidal Activity Of Essential Oil From *Lippia javanica* against *Rhipicephalus appendiculatus* larvae

Caroline J. Kosgei^{1*}, Josphat C. Matasyoh², Charles M. Mwendia¹,

¹Department of Biochemistry, Egerton University. P. O. Box 536-20115 Egerton Kenya

²Department of Chemistry, Egerton University. P. O. Box 536-2011 Egerton, Kenya.

Abstract: Animal ectoparasites particularly ticks contribute majorly to decreased productivity in the livestock industry globally, due to diseases they transmit. This study evaluated the larvicidal activity of essential oil of *Lippia javanica* leaves against larvae of *Rhipicephalus appendiculatus*. The oil was obtained by hydro-distillation of fresh leaves and oil components analysed using Gas Chromatography-Mass Spectrometry (GC-MS). Major components in the oil were monoterpenes as shown by the high percentage of Artemisia ketone (49.52%), m-tert-Butylphenol (8.73%), Linalool (4.43%), beta-myrcene (3.13%), Targetone (2.85%) and Isopiperitenone (2.13%). Sesquiterpenes constituted only 6.06% of the total oil. Contact toxicity was used in the bioassay and mortality data obtained at 6, 12, 24 and 48 hrs. The data obtained during bioassay was then subjected to probit regression analysis to estimate concentration dependent mortality for LC₅₀ and LC₉₀ values in mg/ml. The LC₅₀ in mg/ml were: 3.1(3.0-3.2), 3.1(3.0-3.2) 3.1 (3.0-3.2) 3.07(2.94-3.19) and LC₉₀ were 3.9(3.7-4.2), 3.9(3.7-4.2), 3.9(3.7-4.2), 3.9(3.7-4.2) at 6, 12, 24 and 48 hrs respectively from the start of the experiment. The results of one way ANOVA showed there was no significant difference in activity of the oil against the larvae between 6, 12, 24 and 48 hrs in all the concentrations of the oil used with P>0.05. The findings showed that essential oil of *L. javanica* possessed larvicidal properties and can be used to control tick larvae.

Key words: Cytotoxicity, Phytochemicals, Bioassay

Date of Submission: 01-12-2017

Date of acceptance: 30-12-2017

I. Introduction

Ticks are blood-sucking ectoparasites majorly contributing to decreased productivity in the livestock industry globally due to diseases they transmit. They transmit diseases such as *anaplasmosis*, *babesiosis*, *theileriosis*, tick paralysis and *ehrlichiosis* among others (Piesman and Eisen, 2008). Ticks belong to phylum Arthropoda, class Arachnida and subclass Acari. It is a cosmopolitan group with an estimated 50,000 described tick species (Halliday et al., 2000). Among tick species, *R. appendiculatus* is of major economic importance in Africa since it transmits *Theileria parva*, which causes *theileriosis* a threatening disease in the livestock industry in Eastern, Southeastern and Central Africa. It also transmits corridor disease in cattle, Nairobi sheep disease virus and thogoto virus to livestock (Walker et al., 2000). Synthetic acaricidal agents currently used are associated with resistance, high costs, presence of chemical residues in animal products and environmental pollution. The concerns about negative effects associated with current acaricidal agents drive efforts in bioprospecting for novel compounds that possess acaricidal activity and are safe. Essential oils from plants have been reported to have various bio-efficacies, which include; antibacterial, antifungal, antiviral, antiparasitic and insecticidal (Plaza et al., 2004). The enormous bioactivity of the essential oils has been attributed to low molecular weight of the volatile components present in the oil enabling them to diffuse rapidly through the skin membranes and thus reaching target sites quickly (Gutierrez et al., 2009). Among plant essential oils that have been reported to exhibit acaricidal properties is the essential oil from *Eucalyptus citriodora* which achieved 85% mortality against *Dermanyssus gallinae* mite (David et al., 2009). Similarly, essential oil from *Origanum onites* exhibited acaricidal activity against *Rhipicephalus sturanicus* (Coskun et al., 2008) while essential oil of *Lippia gracilis* Schauer showed activity against *Rhipicephalus (Boophilus) microplus* (de Oliveira et al., 2013). The plant *L. javanica* belongs to *Verbenaceae* family that has about 200 species. These plants are often aromatic due to presence of essential oils and are distributed throughout Central and South Americas as well as in tropical Africa (Arthur et al., 2011). Previous studies on *L. javanica* essential oil showed it had strong and lasting repellent activity against adult *Hyalomma marginatum* cattle tick (Magano et al., 2011). *L. javanica* aqueous leaf extracts at 10% and 20% w/v were found to be effective at controlling *R. appendiculatus*, *R. evertsi*, *B. decoloratus* and *Hyalomma* species (bont-legged ticks) found at study site in Zimbabwe (Madzime et al., 2011). Tea infusions of the leaves of *L. javanica* are used against common symptoms of HIV and AIDS, treatment of lung infections, dysentery and diarrhea (Palgrave et al., 2003). The aim of this study was thus to determine the chemical components present in *L. javanica* oil and its larvicidal properties against *R. appendiculatus*.

II. Materials And Methods

Sample collection

Leaves of *L. javanica* were collected from botanical-garden of Egerton University in Kenya which is at an altitude of 2127 M. A voucher specimen has been deposited at the Department of Biological Sciences, Egerton University.

Extraction of essential oil

Fresh leaves of *L. javanica* were subjected to hydro-distillation in a modified clevenger-type apparatus for at least 4 hrs according to British pharmacopoeia (Papachristos and Stamopoulos, 2004). The essential oil obtained was dried over anhydrous sodium sulphate then stored in sealed glass vials at 4°C awaiting chemical composition analysis using GC-MS and larvicidal bioassay done.

GC-MS Analysis

The extracted essential oil was first diluted with methyl-t-butyl ether (MTBE) (1:100) and analyzed on Agilent GC-MSD apparatus that was outfitted with a Rtx-5SIL MS ('Restek') (30 mm × 0.25 mm, 0.25 µm film thickness). The GC column temperature was maintained at 50°C for 2 min, and then programmed at a temperature of 260°C at the rate of 5°C/min and was held at 260°C for 10 min. Helium was used as the carrier gas (at 0.8 ml/min). The samples were injected in split mode at a ratio of 1:10 to 1:100. The temperature was maintained at 250°C while the transfer line was kept at the temperature of 280°C. The operation of MS was done in the electron impact ionization (EI) mode set at 70 eV, in m/z that ranged from 42 to 350. Compound identification was achieved by comparing the mass spectra and retention indices with ones from the literature (Adams, 2007). It was also supplemented by Wiley and QuadLib 1607 GC-MS libraries. The relative proportions of the essential oil constituents were expressed as percentages obtained by peak area normalization while all relative response factors being taken as one.

Larval bioassay

The larvae used for the bioassay were reared according to Bailey (1960). The oil was first dissolved in 2% Dimethyl Sulfoxide (DMSO) and a stock solution of 4 mg/ml made. The stock solution was obtained after conducting preliminary screening. Serial dilution of 4 mg/ml resulted in 12 concentrations ranging from 1.5 to 4 mg/ml. The bioassay was done using contact toxicity according to Pamoet *al.*, (2005) with modification. This involved attaching Whatman No. 1 filter paper to the bottom of the Petri dish (15 cm) using double sided cellophane tape and 20 larvae placed inside. Different concentrations of the oil was then sprinkled on the Petri dish containing the larvae using a pasture pipette making sure that filter paper was wet and the larvae came in contact with the sprinkled extract. The Petri dish was kept at 25 °C and 75% relative humidity and experiment replicated three times. From the start of the experiment, mortality data was obtained at 6, 12, 24, and 48 hrs under a microscope x 4.5. The larvae were considered dead if they do not move their appendages when prodded with a pin. A negative control was set consisting of 2% DMSO while positive control consisted of 0.2% v/v of amitraz®.

III. Results

Table 1 shows the results of the GC-MS analysis of *L. javanica* essential oil. Major compounds were monoterpenes and these were artemisia ketone (49.52%), m-tert-Butylphenol (8.73%), Linalool (4.43%), beta-myrcene (3.13%), Targetone (2.85%) and Isopiperitenone (2.13%). Table 2 shows mean percentage larval mortalities obtained at 6, 12, 24, and 48 hrs after treatment. LC₅₀ in mg/ml were: 3.1(3.0-3.2), 3.1(3.0-3.2), 3.1(3.0-3.2), 3.1(2.9-3.2) and LC₉₀ were 3.9(3.7-4.2), 3.9(3.7-4.2), 3.9(3.7-4.2), 3.9(3.7-4.2) at 6, 12, 24 and 48 hrs respectively from the start of the experiment. The negative control used was 2% of DMSO which showed no activity against the larvae within 0 to 48 hrs. The results of one way ANOVA showed there was no significant difference in activity of the oil against the larvae between 6, 12, 24 and 48 hrs in all the concentrations of the oil used with P>0.05.

Table 1: GC-MS analysis of *L. javanica* essential oil

S/N	R.T (min)	Compound name	% Concentration	Detection method
1	8.45	β - Myrcene	3.13	GC/MS
2	9.18	Menthatriene	0.51	GC/MS
3	10.12	Targetone	2.85	GC/MS
4	11.23	Bicyclo[3.1.0]hexane,6-ethylene-	0.44	GC/MS
5	11.46	Linalool	4.43	GC/MS
6	12.67	Camphor	0.90	GC/MS
7	13.11	Artemisia Ketone A21	49.52	GC/MS
8	15.21	Phenol, M-tert butyl (Ocimenone)	8.73	GC/MS
9	16.28	Isopiperitenone	2.13	GC/MS

10	20.06	Z-caryophyllene	1.99	GC/MS
11	20.91	4,7,10-Cycloundecatriene,1,1,4,8-ctramethyl-, cis, cis, cis-	0.92	GC/MS
12	24.05	Caryophellene oxide	1.31	GC/MS

Table 2: Mean percentage larval mortalities obtained at 6, 12, 24, and 48 hrs

Concentration in mg/ml	Mean % larval mortality at the hours shown below			
	0-6	6-12	12-24	24-48
1.50	0±0	0±0	0±0	0±0
2.00	0±0	0±0	1.7±2.9	5±5
2.22	8.3±7.6	8.3±7.6	8.3±7.6	10±5
2.50	13.3±5.8	15±5	15±5	16.7±2.9
2.80	28.3±5.8	33.3±2.9	36.7±2.9	38.3±2.9
3.00	38.3±2.9	38.3±2.9	43.3±2.9	48.3±2.9
3.20	50±8.7	51.7±5.8	56.7±7.6	60±5
3.45	61.7±7.6	63.3±7.6	66.7±7.6	73.3±5.8
3.65	70±5	71.7±7.6	76.7±2.9	80±5
3.70	85±5	86.7±2.9	90±5	96.7±5.8
3.85	96.7±2.9	100±0	100±0	100±0
4.00	100±0	100±0	100±0	100±0
Amitraz @ (0.2 % v/v) ^P	0±0	53.3±5.8	90±10	100±0
DMSO+distilled H ₂ O) ^Q	0±0	0±0	0±0	0±0

^Ppositivecontrol, ^QNegativecontrol

IV. Discussion

Major essential oil components present were monoterpenes with percentage yield of 70.79. This compares with essential oil of *Lippiapalmeri* which showed monoterpenes (70.35%) being dominant (Ortega-Nieblas *et al.*, 2011). Also essential oil of *Vitexagnus-castus* L was dominated by high amount of monoterpenes and it constituted the predominant chemical class in the oils from the leaves and fruit. The similarity in dominance could be because the plants are of the same genus thus similar chemotype. Nonetheless, variation in essential oil components have been observed even within the same species. Variability in quantity and composition of essential oil in plants have been attributed to their developmental stage, climate factors, altitude, harvest season and plant handling (Dunford and Silva, 2005). The dominance of myrcene and linalool in this study compared with *L. javanica* oil in western Zimbabwe which showed the oil majorly contained myrcene (54.0 %) and linalool (68.8 %) (Chagonda and Chalchat, 2015), with the difference in concentration being brought about by difference in ecological conditions. Though Artemisia Ketone reported the highest concentration in this study (49.52), previous studies in Zimbabwe showed the plant had trace amount of this component i.e less than 0.05% indicating difference in climatic conditions (Chagonda and Chalchat, 2015). There is similarity in the major compounds observed with that of yegonet *al.*, (2015). This is because the plant was obtained from the same geographical locality.

The activity observed in the oil against the tick larvae could be due to synergistic action of major components present in the oil, whose acaricidal bioefficacy have been determined (Park *et al.*, 2008). Nevertheless, minor components present in the oil may also have contributed to the biological activity of the oil (Iacobellis *et al.*, 2005). Among the major compounds present in the oil whose bio-activity has been determined include β-myrcene, tagetone, linalool, camphor, and artemesia ketone.

Myrcene has been found to be repellent against *R. appendiculatus* and *Sitophilus zeamais* Motschulsky (Ndunguet *al.*, 1995). Linalool obtained from *ocimum canum* has been reported to be toxic against *Zabratessubfasciatus* a coleopteran (Weaver *et al.*, 1991) while Linalool isolated from molasses grass *Melinis minutiflora* was found to be lethal to 100 % of *R. microplus* tick larvae in 15 min (Prates *et al.*, 1998). Commercial camphor and terpinen-4-ol were found to be 7 to 48 times more toxic against both male and female *A. obtectus* adults' mite than the less active monoterpenes tested (Papachristos *et al.*, 2004). Tagetone was one of the major compound in *Tagetes minuta* essential oil that had over 95% efficacy against four species of ticks at a concentration of 20% (Garcia *et al.*, 2012).

From the results of the bioassay, there was no significant difference in activity between 6, 12, 24 and 48 hrs in *L. javanica* with $P > 0.05$. This is also similar to what was observed in *L. kituiensis* when it was exposed to the same larvae of *R. appendiculatus* (Kosgei *et al.*, 2014). This explains the fact that essential oils are volatile compounds, which are likely to evaporate, and are prone to rapid aerial oxidation and chemical re-arrangement hence loss of activity quickly. Therefore, further increase in time could not increase larval mortality (Birkett *et al.*, 2008).

Results of this study indicate that essential oils obtained from *L. javanica* plant have acaricidal properties and could be used as lead molecules for formulation of acaricidal drugs against larvae of *R. appendiculatus*.

Acknowledgements

The authors are grateful to Egerton University for availing their research laboratories for conducting the research. The authors are also thankful to Richard Ochieng for helping in the rearing of *R. appendiculatus* larvae.

Reference

- [1]. Adams, R. P. (2007). Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry. 4th edn. Allured publishing corporation: Carol Stream, Illinois, USA.
- [2]. Arthur, H., Joubert, E., De Beer, D., Malherbe, C. J., and Witthuhn, R. C. (2011). Phenylethanoid glycosides as major antioxidants in Lippia multiflora herbal infusion and their stability during steam pasteurization of plant material. *Journal of Food Chemistry*, **127**: 581–588.
- [3]. Bailey K. P. 1960 — Note on the rearing of *Rhipicephalus appendiculatus* and their infection with *Theileriaparva* for experimental transmission. *Bulletin of epizootic diseases of Africa*, **8**: 33-43.
- [4]. Chagonda, L. S. and Chalchat, J. (2015). Essential oil Composition of *Lippia javanica* (Burm.f.) Spreng chemotype from Western Zimbabwe. *Journal of Essential Oil Bearing Plants*, **18**(2): 482-485.
- [5]. Coskun, S., Girisgin, O., Kürkcüoğlu, M., Malyer, H., Girisgin, A. O., Kirimer, N., Baser, K. H. (2008). Acaricidal efficacy of *Origanum monites* L. essential oil against *Rhipicephalus turanicus* (Ixodidae). *Parasitol Research*, **103**: 259-261.
- [6]. David, R. G., Dino, M., Olivier, A. E. S., Jonathan, H. G. (2009). Variation in chemical composition and acaricidal activity against *Dermatophylos gallinae* of four eucalyptus essential oils. *Experimental and Applied Acarology*, **48**: 43–50
- [7]. de Oliveira C. E. M., Costa-Junior, L. M., Pinto, J. A. O., De Alexandria Santos, D., De Araujo, S. A., De Fátima Arrigoni-Blank, M., and Blank, A. F. (2013). Acaricidal Activity of *Lippia gracilis* Essential Oil And Its Major Constituents. *Rhipicephalus (Boophilus) Microplus. Veterinary Parasitology*, **195**(1): 198-202.
- [8]. Dunford N T, V R Silva (2005) Effect of water stress on plant growth and thymol and carvacrol concentration in Mexican oregano grown under controlled conditions. *J. Appl. Horti.* **7**: 20-22.
- [9]. Garcia, M. V., Matias, J., Barros, J. C., Lima, D. P. D., Lopes, R. D. S., & Andreotti, R. (2012). Chemical identification of *Tagetes minuta* Linnaeus (Asteraceae) essential oil and its acaricidal effect on ticks. *Revista Brasileira de Parasitologia Veterinária*, **21**(4): 405-411.
- [10]. Gutierrez J, Bourke P, Lonchamp J. (2009). Impact of plant essential oil on microbiological and quality markers of minimally processed vegetables. *Food Science and Environmental Health*, **10** (2): 195–202.
- [11]. Halliday, R. B., Oconnor, B. M. and Baker, A. S. (2000). Nature and human society: The quest for a sustainable world. In P. H. Raven and T. Williams (eds.), *Global diversity of mites* pp 192–203. Washington, D C: National Academy Press.
- [12]. Iacobellis, N. S., Lo Cantore, P., Capasso, F., Senatore, F. (2005). Antibacterial activity of *Cuminum cyminum* and *Carum carvi* essential oils. *Journal of Agricultural Food Chemistry*, **53**: 57–61.
- [13]. Kosgei, C. J., Matasyoh, J. C., Mwendia, C. M. (2014). Chemical composition and larvicidal activity of essential oil of *Lippiakituiensis* against larvae of *Rhipicephalus appendiculatus*. *Int. J. Biol. Chem. Sci.*, **8**(4): 1938-1947.
- [14]. Madzimure, J., Nyahangare, E. T., Hamudikuwanda, H., Hove, T., Stevenson, P. C., Belmain, S. R., Mvumi, B. M. (2011). Acaricidal efficacy against cattle ticks and acute oral toxicity of *Lippia javanica* (Burm. F.) Spreng. *Tropical Animal Health Production*, **43**: 481–489.
- [15]. Magano, S. R., Nchu, F., Eloff, J. N. (2011). In vitro investigation of the repellent effects of the essential oil of *Lippia javanica* on adults of *Hyalomma marginatum*. *African Journal of Biotechnology*, **10**(44): 8970-8975.
- [16]. Ndungu, M., Lwande, W., Hassanali, A., Moreka, L., and Chhabra, S. C. (1995b). *Cleome monophylla* essential oil and its constituents as tick (*Rhipicephalus appendiculatus*) and maize weevil (*Sitophilus zeamais*) repellents. *Entomologia Experimentalis et Applicata*, **76**: 217-222
- [17]. Ortega-Nieblas, M. M., Robles-Burgueño, M. R., Acedo-Félix, E., González-León, A., Morales-Trejo, A., & Vázquez-Moreno, L. (2011). Chemical composition and antimicrobial activity of oregano (*Lippia palmeri* S. WATS) essential oil. *Revista Fitotecnia Mexicana*, **34**(1): 11-17
- [18]. Palgrave, M. C., Drummond, R. B. and Moll, E. J. (2003). *Trees of southern Africa*. 3rd Edition, Cape Town: Struik.
- [19]. Pamo, E. T., Tendongeng, F., Kana, J. R., Payne, V. K., Boukila, B., Lemoufouet, J., Miegoue, E., Nanda, A. S. (2005). A study of the acaricidal properties of an essential oil extracted from the leaves of *Ageratum houstonianum*. *Vet. Parasitol.*, **128**: 319-323
- [20]. Papachristos, D. P. and Stamopoulos, D. C. (2004). Fumigant toxicity of three essential oils on the eggs of *Acanthoscelides obtectus* Say (Coleoptera: Bruchidae). *Journal of Stored Products Research*, **40**: 517-525.
- [21]. Papachristos, D. P., Karamanoli, K. I., Stamopoulos, D. C., Menkissoglu-Spiroudi, U. (2004). The relationship between the chemical composition of three essential oils and their insecticidal activity against *Acanthoscelides obtectus*. *Pest Management Science*, **60**: 514–520.
- [22]. Park, I. K., Kim, J. N., Lee, Y. S., Lee, S. G., Ahn, Y. J., Shin, S., C. (2008) Toxicity of plant essential oils and their components against *Lycoriella ingenua* (Diptera: Sciaridae). *Journal of Economic Entomology*, **39**: 275–279.
- [23]. Piesman, J. and Eisen, L. (2008). Prevention of tick-borne diseases. *Annual Review of Entomology*, **53**: 323-343.
- [24]. Plaza, P., Torres, R., Usall, J., Larmaca, N and Viñasi, I. (2004). Evaluation of the potential of commercial postharvest application of essential oils to control citrus decay. *Journal of Horticultural Science and Biotechnology*, **79** (6): 935-940.
- [25]. Prates, H. T., Leite, R. C., Craveiro, A. A., Oliveira, A. B. (1998). Identification of some chemical components of the essential oil from molasses grass (*Melinis minutiflora* Beauv.) and their activity against cattle tick (*Boophilus microplus*). *Journal Brazilian Chemistry Society*, **9**: 193–197.
- [26]. Walker, J. B., Keirans, J. E. and Horak, I. G. (2000). *The Genus Rhipicephalus*. Cambridge, Cambridge University Press.
- [27]. Weaver, D. K., Dunkel, F. Y., Ntezurbanza, L., Jackson, L. L Stock, D. T. (1991). The efficacy of linalool a major component of freshly *Ocimum canum* Sims (Lamiaceae) for protection against post harvest damage by certain stored coleopteran. *Journal of Stored Products Research*, **22**: 213-220.
- [28]. Yegon, K. P., Matasyoh, J. C., Wagara, I. N. (2015). Chemical composition and antifungal activity of the essential oil from *Lippia javanica* (Verbenaceae). *International Journal of Biotechnology and Food Science*, **4**(1):1-6.

Caroline J. Kosgei "Chemical Composition And Larvicidal Activity Of Essential Oil From *Lippia javanica* against *rhipicephalus appendiculatus* larvae." *IOSR Journal of Pharmacy and Biological Sciences (IOSR-JPBS)* **12.6** (2017): 46-49.