

## Reproductive Organ Damage of Domestic Ruminants Caused By Animal African Trypanosomiasis: A Review

Abdullahi, M. A.<sup>1\*</sup>, Hassan, A.<sup>2</sup>, Ibrahim, M. L.<sup>3</sup>, Ubale, C.<sup>4</sup>, Musa, T. I.<sup>4</sup>

<sup>1</sup>Trypanosomiasis Research Department, Nigerian Institute for Trypanosomiasis (and Onchocerciasis) Research, No. 1 Surame road, Ungwan rimi GRA, Kaduna, Kaduna State, Nigeria.

<sup>2</sup>Sokoto State Liaison Office, Nigerian Institute for Trypanosomiasis (and Onchocerciasis) Research, Sokoto State.

<sup>3</sup>Biotechnology Unit, Nigerian Institute for Trypanosomiasis (and Onchocerciasis) Research, No. 1 Surame road, Ungwan rimi GRA, Kaduna, Kaduna State, Nigeria.

<sup>4</sup>Vector and parasitological Studies Research Department, Nigerian Institute for Trypanosomiasis (and Onchocerciasis) Research, No. 1 Surame road, Ungwan rimi GRA, Kaduna, Kaduna State, Nigeria.

\*Corresponding author: E-mail: goodthinking338@gmail.com

---

**Abstract:** Reproductive disorders have been recognized in animal African trypanosomiasis since 19<sup>th</sup> century. The damages caused by trypanosomes have direct or indirect link to some specific organs like endocrine and pituitary gland which secretes follicle stimulating hormone (FSH), interstitial-cell stimulating hormone (ICSH) and growth hormone (GH). These hormones play an important role in the spermatogenic cycle in males and oestrus cycle in females. However, the pathogenesis of these damages has not been clearly elucidated. We believe that this aspect deserves closer study especially in livestock where reproductive performance is the cornerstone of productivity. It is hoped that this review which is centered on the chronological report of works that has been carried out on some reproductive organs damage in selected domestic ruminants infected with trypanosomes will add more light to the pathophysiology of these organs.

**Keywords:** Trypanosomiasis, pituitary gland, endocrine, ruminant, spermatogenic and oestrus.

---

Date of Submission: 02-02-2017

Date of acceptance: 07-12-2017

---

### I. Introduction

Trypanosomiasis is a disease peculiar to sub-Saharan African which affect both human (sleeping sickness) and animal (Nagana or Samore' in cattle). African animal trypanosomiasis is a disease affecting virtually every organ thereby causes anaemia at the first stage of infection and death on the long run (Rodrigues *et al.*, 2012). Trypanosome is transmitted cyclically through biting of tsetse fly (*Glossina spp.*) and remains one of the major infections limiting the animal growth and livestock industry in Africa (OIE, 2013). *Trypanosoma brucei*, *T. congolense*, and *T. vivax* are the major pathogenic trypanosomes peculiar to animals (Desquesnes, 2004). Trypanosomiasis is characterized by anaemia, anorexia, depression, increased respiratory and heart rates, nasal discharge, subcutaneous oedema, intermittent fever, enlargement of lymph nodes, central nervous system and reproductive disorders (Bentivoglio *et al.*, 1994; Desquesnes, 2004). Considering trypanosome effect on reproductive disorders in male ruminants, reports have shown that it can cause severe testicular degeneration, scrotal inflammation, penile protrusion, prepuccial inflammation, epididymitis, abnormal spermatogenesis and deterioration of semen characteristics (Sekoni, *et al.*, 1990; Adamu *et al.*, 2007; Victor *et al.*, 2012).

In female ruminants, Faye *et al.* (2004) and Silva *et al.* (2013) documented that the disease causes irregular oestrus cycle, cystic degeneration of the ovary, follicular cyst, decreased conception rate, abortion, low birth weight and neonatal death. In view of these, an estimated 50 million heads of cattle are at risk, and in a situation where the disease is tolerable, up to 50% cases of mortality and morbidity are recorded in animals and consume a sum of \$600million to \$1.2billion annually in Africa (FAO, 1994; Maudlin, 2006). This threat has been noticed to hinder food availability (Samdi *et al.*, 2010).

Reproductive disorders have been recognized in animal African trypanosomiasis since the early part of 19<sup>th</sup> century (Ikede *et al.*, 1988). We believe that this aspect deserves closer study especially in livestock where reproductive performance is the cornerstone of productivity. It is hoped that this review which is centered on the chronological report of works that has been carried out on some reproductive organs damage in selected domestic ruminants infected with trypanosomiasis will add more light to the pathophysiology of these organs.

## 2.1 THE ENDOCRINE SYSTEM

The endocrine system plays a central role in the regulation of important body functions such as growth, differentiation, reproduction, maintenance of the internal environment, and adaptation to changes in the external environment. Its involvement in reproduction has direct link to hypothalamus and anterior pituitary (Senger, 2005). The gonadotropin releasing hormone (GnRH) produced from the hypothalamus induces release of gonadotropin, follicle stimulating hormone (FSH) and luteinising hormone (LH) from the anterior pituitary. FSH and LH are required for spermatogenesis and sperm maturation, while development of male secondary characteristics and libido depends on the testosterone (Khisk, 2008). *T. congolense* and *T. vivax* are known to be intravascular and non-tissue invasive which is capable of affecting endocrine system. The mechanism of these endocrine lesions has been postulated on the ability of the trypanosomes to localize in the organs thereby causing severe damage (Raheem, 2014).

## 2.2 PITUITARY GLAND

Pituitary hormones (gonadotropins) play a major role in the reproductive process. In particular, follicle stimulating hormone (FSH), interstitial-cell stimulating hormone (ICSH) and growth hormone (GH) play an important part in the spermatogenic cycle in males and oestrus cycle in females. Consequently, absence of pituitary gonadotropins leads to a block in sperm maturation in rats (Clermont and Morgentaler, 1955) or to marked testicular degeneration and abortion in man and animals (Steinberger and Duckett, 1967; Apter, 1970; Ikede and Losos, 1975). The pituitary gland, which is connected to the central nervous system (CNS) through the hypothalamus, is one of the endocrine organs affected by trypanosomiasis. Ikede and Losos (1979) reported focal coagulative necrosis and interstitial mononuclear infiltration in pituitaries of sheep experimentally infected with *T. brucei*, these lesions were associated with extra-vascular localization of trypanosomes in the pituitary gland. Studies on infected domestic and laboratory animals have provided more evidence of specific damage to the pituitary gland (Ikede *et al.*, 1988). Extensive mononuclear inflammation of the gland has been described in cattle, sheep and goats experimentally infected with *T. brucei* (Losos and Ikede, 1970, 1972; Ikede *et al.*, 1977; Moulton and Sollod, 1976; Morrison *et al.*, 1981). Abebe *et al.* (1993) reported that the vessels of the microvasculature in the pituitary glands of infected animals with *T. Congolense* were highly distended with trypanosomes, erythrocytes, macrophages, cellular debris, pituitary cell secretory granules, and microvasculature dilation in skeletal muscle, myocardium, and brain. These effects directly reduce the availability of necessary hormones required during mating; thus hindering normal reproductive processes in ruminants. Extensive dilation of the pituitary microvasculature could lead to pooling of blood in the capillary beds, affecting the efficiency of circulation through the capillaries and in turn possibly causing the impairment of normal nutrient and metabolite exchange, leading to cellular apoptosis (Losos and Ikede, 1972; Wellde *et al.*, 1989; Nyeko *et al.*, 1990). Reincke *et al.* (1998) reported that animal African trypanosomiasis caused local inflammation of the pituitary and the gonads, associated with increase in the level of cytokines such as tissue necrotic factor  $\alpha$  (TNF-  $\alpha$ ) and interleukins 6 (IL-6). Studies on hormonal assay under trypanosomiasis infection revealed decline in the level of Luteinsing hormone and testosterone in ruminants (Waindi *et al.*, 1986; Boly *et al.*, 1994; Mutayoba *et al.*, 1994).

## 2.3 FEMALE REPRODUCTIVE ORGANS

Ovary is the primary female reproductive organ and has two important functions: producing the female reproductive cell (the egg or ovum) and producing the hormones estrogen and progesterone. The secondary sex organs are a series of tubes that receive semen, transport sperm to the egg so it can be fertilized, nourish the fertilized egg (embryo), and allow the calf to be birthed. These organs include the vagina, cervix, uterus, uterine horns, and oviducts (also called Fallopian tubes), which each has a funnel-shaped opening called the infundibulum. The released egg is caught by the infundibulum and moves into the oviduct, where fertilization occurs if viable sperm are present. The egg remains capable of fertilization for only a few hours. Thus, it is very important that fertile sperm be present near the time of ovulation. The egg moves through the oviduct and into the uterus within the next three to four days. Following the trend of embryological development, any alteration may terminate the process and thereby causes infertility in ruminants.

Histopathological examination of *Trypanosoma brucei* infected West African dwarf does by Leigh *et al.* (2014) revealed subacute necrotising adenohypophysitis characterized by widespread necrosis and disruption of the architecture of the adenohypophysis. Multiple foci of inflammatory cells which were mostly lymphocytes and a few neutrophils were also observed in the pituitary gland. Also, the hypothalamus of the infected does showed widespread congestion of blood vessels with severe perivascular lymphocytic cuffs meanwhile uterus showed acute severe endometritis characterized by marked extensive necrosis of the epithelium with varying degrees of necrosis and distortion of the tubular glands in the endometrial stroma. Acute placentitis characterized by marked vascular congestion and multifocal lymphocytic aggregates was also observed in infected pregnant does. The lesion observed in the uterine body, a severe necrotising endometritis, is capable of

reducing its potential to carry out its physiological functions which include the production of prostaglandin F<sub>2α</sub> (i.e. luteolysin) and nidation (Leigh and Fayemi, 2013). *T. vivax* causes lesions in the placenta of ewe (Silva *et al.*, 2013). These lesions are capable of interfering with normal exchanges between foetus and dam, leading to foetal hypoxia and death *in utero* or the initiation of early dehiscence of the placenta and subsequent abortion (Dubey *et al.*, 2006). *T. brucei* possesses the ability to traverse the placenta causing placentitis and also infection *in utero* which may be linked to cord atrophy due to impaired nutritional and gaseous exchanges between the dams (Marcato *et al.*, 1991; Rocha *et al.*, 2004). Adenowo *et al.* (2005) observed inflammatory responses of hypothalamus and adenohipophysis when examining histological changes in *T. vivax* infected sheep. Leigh and Fayemi (2013) reported reduction in the concentrations of follicle stimulating hormone during oestrus, and luteinizing hormone during proestrus and oestrus, as well as oestradiol during the follicular phase in *T. brucei* infected West African Dwarf does.

## 2.4 MALE REPRODUCTIVE ORGANS

Trypanosome infection has been reported to cause reproductive dysfunction both in male and female ruminants. In males, changes in semen characteristics and testicular pathology are among the reported abnormalities observed in trypanosome infection. Inflammatory changes in the genital organs are usually mild or absent but there is progressive and marked testicular degeneration that can lead to atrophy and aspermia (Isoun and Anosa, 1974; Isoun *et al.*, 1975; Anosa and Isoun, 1980; Kaaya and Oduor-Okelo, 1980; Masake, 1980; Anosa, 1983). The cause of the testicular lesion is believed to be due to the effect of prolonged fever, thrombosis of spermatic blood vessels, and the general wasting of body organs (Anosa and Isoun, 1980; Anosa, 1983). In male animals infected with the *brucei* group, the lesions are a combination of scrotal dermatitis, orchitis and periorchitis. Soon after infection the parasites localize in the scrotal skin and hydrocoel fluid and also invade the tunica vaginalis, testis, epididymis and spermatic cord, provoking a non-purulent granulomatous inflammation in sheep (Ikede, 1979).

The severity of testicular and epididymal lesions is reflected in poor quality of semen and the high percentage of abnormal spermatozoa present in the ejaculate of bulls (Isoun *et al.*, 1975; Grundler, 1985) and rams (Isoun and Anosa, 1974; Anosa and Isoun, 1980; Akpavie *et al.*, 1987) experimentally infected with *T. vivax*, *T. congolense* or *T. brucei*. It has also been shown in rabbits and sheep that successful treatment with trypanocides will lead gradually to normal spermatogenesis over a period of several months if the original lesions have not been complicated by secondary bacterial infections (Ikede and Akpavie, 1982; Akpavie *et al.*, 1987).

Sekoni *et al.* (1990) revealed the scrotal oedema and testicular degeneration characterized by degenerated spermatogenic cells in bull infected with *T. congolense*. Scrotal oedema can also occur as a result of inflammation caused by the parasite (Rakha *et al.*, 2006). Okubanjo *et al.* (2014) reported reduction in semen concentration and increase in percentage sperm abnormalities of Yankasa rams infected with *T. congolense*. Low semen concentrate was also observed in rams infected with *T. brucei* and *T. vivax* (Sekoni, 1992) which may be attributed to localization of trypanosomes in the scrotal skin, provoking non-purulent inflammation that leads to degeneration of the seminiferous and spermatozoa abnormalities.

## II. Conclusion

In Trypanosomiasis infection, the animal becomes less productive because of the damages done to the reproductive organs by the trypanosome parasites. These damages are either directly to the organs or to the microvascular structures of the organs. The pathogenesis of these damages has not been clearly elucidated. We believe that this aspect deserves closer study especially in livestock where reproductive performance is the cornerstone of productivity.

## References

- [1]. Abebe, G. R. Eley, R. M. and Kole-Moiyoi, O. (1993). Reduced responsiveness of the hypothalamic-pituitary-gonadal axis in Boran cattle infected with *T. congolense*. *Acta Endocrinology (Copenhagen)*, 129:74-80
- [2]. Adamu, S., Fatihu, M. Y., Useh, N. M., Mamman, M., Sekoni, V.O. and Esievo, K. A. N. (2007). Sequential testicular and epididymal damage in Zebu bulls experimentally infected with *Trypanosoma vivax*. *Veterinary Parasitology*, 143(1): 29-34
- [3]. Adenowo, T. K., Njoku, C. O., Oyedipe, E. O. and Sanusi, A. (2005). Lesion of the hypothalamus, adenohipophysis and the ovaries in *Trypanosoma vivax* infected Yankasa Ewes. *Nigerian Veterinary Journal*, 26:56-62
- [4]. Akpavie, S. O., Ikede, B. O. and Egbunike, G. N. (1987). Ejaculate characteristic of sheep infected with *Trypanosoma brucei* and *Trypanosoma vivax*: Changes caused by treatment with Diminazine aceturate. *Research in Veterinary Science*, 42:1-6
- [5]. Anosa, V. O. (1993). Diseases produced by *Trypanosoma vivax* in ruminants, horses and rodents. *Veterinary Medicine*. B 30:111-141
- [6]. Anosa, V. O. and Isoun, T. T. (1980). Further observation on the testicular pathology in *Trypanosoma vivax* infection of sheep and goats. *Research in Veterinary Science* 28:151-160
- [7]. Apted, P. I. C (1970). Clinical manifestation and diagnosis of sleeping sickness in the African trypanosomiasis, Reviewed by H. W. Mulligan and W. H. Potts. *Allen and Unwin, London*. Pg681-683

- [8]. Bentivoglio, M., Grassi-Zucconi, G., Olsson, T. and Kristensson, K. (1994). *Trypanosoma brucei* and the nervous system. *Trends in Neuroscience*, 17(8): 325-329
- [9]. Boly, H., Humblot, P., Tillet, Y. and Thibier, M. (1994). Effects of *Trypanosoma congolense* infection on the pituitary gland of Baoule bulls: Immunohistochemistry of LH- and FSH-secreting cells and response of plasma LH and testosterone to combined dexamethasone and GnRH treatment. *Journal of Reproduction and Fertility*, 100:157-162
- [10]. Clermont, Y. and Morgentaler, H. (1955). Quantitative study of spermatogenesis in the hypophysectomized rat. *Endocrinology*, 57:369-382
- [11]. Desquesnes, M. (2004). Livestock trypanosomes and their Vectors in Latin America (English translation). Published by *World Organization for Animal Health (OIE)*, pp: 8-19
- [12]. Dubey, J. P., Buxton, D. and Wouda, W. (2006). Pathogenesis of bovine neosporosis. *Journal of Comparative Pathology*, 134:267-289
- [13]. Faye, D., Sulon, J., Kane, Y., Beckers, J. F., Leak, S., Kaboret, Y., Sousa, N. M., Losson, B. and Geerts, S. (2004). Effects of an experimental *Trypanosoma congolense* infection on the reproductive performance of West African dwarf goats. *Theriogenology*, 62:1438-1451
- [14]. Food and Agricultural Organization (1994). *Animal production and Health Papers* No 12, Rome, Italy, 1-3
- [15]. Grundler, G. (1985). The influence of trypanosomiasis on sperm quality. *Proceedings of the 18th meeting of the OAU/ISCTR. Harare, Zimbabwe.*
- [16]. Ikede, B. O. (1979). Genital lesions in experimental chronic *Trypanosoma brucei* infection in rams. *Research in Veterinary Sciences*, 26: 145-151
- [17]. Ikede, B. O. and Akpavie, S. O. (1982). Delay in resolution of trypanosome-induced genital lesions in male rabbits infected with *Trypanosoma brucei* and treated with diminazene aceturate. *Research in Veterinary Science*, 32:374-376
- [18]. Ikede, B. O., Elhassan, E. and Akpavie, S. O. (1988). Reproductive disorders in African trypanosomiasis: A review. *Acta Tropica* 45:5-10
- [19]. Ikede, B. O. and Losos, G. J. (1975). Studies on the pathogenesis of *Trypanosoma brucei* infection in sheep: Hypophysical and other endocrine lesions. *Journal of Comparative Pathology*, 85:37-44
- [20]. Ikede, B. O., Akpokodje, J. U., Hill, D. H. and Ajidagba, P. O. (1977). Clinical haematological and pathological studies in donkeys experimentally infected with *Trypanosoma brucei*. *Tropical Animal Health Production*, 9:93-98
- [21]. Isoun, T. T. and Anosa, V. O. (1974). Lesions in the reproductive organs of sheep and goats infected with *T. vivax*. *Tropenmedizin und Parasitologie*, 25:469-476
- [22]. Isoun, T. T., Akpokodje, J. U. and Anosa, V. O. (1975) Testicular changes in white Fulani zebu (Bunaji) cattle experimentally infected with *Trypanosoma vivax*: A preliminary report. *Journal of Nigerian Veterinary Medical Association*, 4:107-108
- [23]. Kaaya, G. P. and Oduor, O. (1980). The effect of *Trypanosoma congolense* infection on the testis and epididymis of goat. *Bulletin Animal Health Production in Africa*, 28:1-5
- [24]. Khisk, W. H. (2008). Interrelationship between ram plasma testosterone level and some semen characteristics. *Slovak Journal of Animal Science*, 41:67-71
- [25]. Leigh, O. O. and O. E. Fayemi, (2013). The effects of experimental *Trypanosoma brucei* infection on hormonal changes during the oestrous cycle, pregnancy and pregnancy outcome in West African dwarf goat does. *Wayamba Journal of Animal Science*, 5:685-695
- [26]. Leigh, O. O., Emikpe, B. O. and Ogunsola, J. O. (2014). Histopathological changes in some reproductive and endocrine organs of *Trypanosoma brucei* infected West African dwarf goat does. *Bulgarian Journal Veterinary Medicine*, ISSN 1311-1477; online at <http://tru.uni-sz.bg/bjvm/bjvm.htm>
- [27]. Losos, G. J. and Ikede, B. O. (1970). Pathology of experimental trypanosomiasis in the albino rat, rabbit, goat and sheep. A preliminary report. *Canada Journal of comparative medicine*, 34:209-212
- [28]. Losos, G. J. and Ikede, B. O. (1972). Review of pathology of disease in domestic and laboratory animals caused by *T. congolense*, *T. vivax*, *T. rhodesiense* and *T. gambiense*. *Veterinary Pathology*, 9: 1-71
- [29]. Marcato, P. S., Benazzi, C., Vecchi, G., Galeotti, M., Salda, D. L., Sarli, G. and Lucidi, P. (1991). Clinical and pathological features of viral haemorrhagic disease of rabbits and the European brown hare syndrome. *Revue Scientifique et Technique de l'Office International des Epizooties*, 10, 371-392
- [30]. Masake R. A. (1980). The pathogenesis of infection with *Trypanosoma vivax* in goats and cattle. *Veterinary Research* 107:551-557
- [31]. Maudlin, I. (2006). Centennial review: African trypanosomiasis. *Annals of Tropical Medicine and Parasitology*, 100:679-701
- [32]. Morrison, W. L., Murray, M., Sayer, P. D. and Preston J. M. (1981). The pathogenesis of experimentally induced *Trypanosoma brucei* infection in the dog's tissue and organ damage. *American Journal Pathology*, 102:168-181
- [33]. Moulton, J. E. and Sollod, A. E. (1976). Clinical, serologic and pathologic changes in calves with experimentally induced *Trypanosoma brucei* infection. *American Journal of Veterinary Research*. 37. 791- 802.
- [34]. Mutayoba, B. M., Eckersall, P. D., Jeffcoate, I. A. Cestnik, V. and Holmes, P. H. (1994). Effect of *Trypanosoma congolense* infection in ram on the pulsative secretion of LH and testosterone and responses to injection of GnRH. *Journal of Reproduction and Fertility*, 102: 425-431
- [35]. Nyeko, J. H. P., ole-MoiYoi, O. K., Majiwa, P. A. O., Otieno, L. H. and Oci ba, P. M. (1990). Characterization of trypanosome isolates from cattle in Uganda using species-specific DNA probes reveals predominance of mixed infections. *Insect Science Applied*, 11:271-280
- [36]. Office of the International Epizootic (2013). Trypanosomosis-tsetse transmitted. *OIE Terrestrial Manual*, pp:1-11.
- [37]. Okubanjo, O. O., Sekoni, V. O., Ajanusi, O. J. and Adeyeye, A. A. (2014). Semen characteristics and reaction time of Yankasa rams experimentally infected with *Trypanosoma congolense*. *Global Veterinaria* 13(3): 297-301
- [38]. Raheem, K. O. (2014). A review of trypanosomosis-induced reproductive dysfunctions in male animals. *Agrosearch*, 14(1):30-38
- [39]. Rakha, E., Puls, F, Saidul, I and Furness, P. (2006). Torsion of the testicular appendix: Importance of associated acute inflammation. *Journal of Clinical Pathology*, 59: 831-834.
- [40]. Reincke, M., Arlt, W., Heppner, C., Petzke, F., Chrousos, G. P. and Allolio, B. (1998). Neuroendocrine dysfunction in African Trypanosomiasis: The role of cytokines. *Annals New York Academic Science*, 840:809-821
- [41]. Rocha, G., Martins, A., Gama, G., Brandao, F. and Atougua, J. (2004). Possible cases of sexual and congenital transmission of sleeping sickness. *The Lancet*, 363, 247
- [42]. Rodrigues M. M., Oliveira, A. V. and Bellio, A. (2012). The immune response to *Trypanosoma cruzi*: Role of Toll-like receptors and perspectives for vaccine development. *Journal of Parasitology Research*; doi:10.1155/2012/507874

- [43]. Samdi, S. M., Abenga, J. N., Attahir, A., Haruna, M. K., Wayo, B. M., Fajinmi, A. O., Sumayin, H. M., Usman, A. O., Hussaina, J. Z., Muhammad, H., Yamap, J. E., Ovbagbedia, R. P. and Abdullahi, R. A. (2010). Impact of trypanosomosis on food security in Nigeria; A Review *International Journal of Animal and Veterinary Advances*, 2:47-50
- [44]. Sekoni, V. O., Njoku, C. O., Kumi-Diaka J. and Saror, D. I. (1990). Pathological changes in male genitalia of cattle infected with *Trypanosoma vivax* and *Trypanosoma congolense*. *British Veterinary Journal*, 146: 175-180
- [45]. Sekoni, V.O. (1992). Effect of *Trypanosoma vivax* on semen characteristics of Yankasa rams. *British Veterinary Journal*, 148:501-506
- [46]. Senger, P. L. (2005). Pathways to pregnancy and parturition. *Pullman: Current Conceptions Inc.* 2nd edition pp. 373
- [47]. Silva, T. M., Olinda, R. G., Rodrigues, C. M., Camara, C. L., Lopes, F. C., Coelho, W. A. (2013). Pathogenesis of reproductive failure induced by *Trypanosoma vivax* in experimentally infected pregnant ewes. *Veterinary Research*, 44: 1.
- [48]. Steinberger, E., Duckett, G. E. (1967). Hormonal control of spermatogenesis. *Journal of Reproductive Fertility*, 2:75-87
- [49]. Victor, I., Sackey, A. K. B. and Natala, A. J. (2012). Penile protrusion with hemorrhages and prepuccial inflammation in pigs experimentally infected with *Trypanosoma congolense*. *Journal of Animal Production Advances*, 2(6): 297-302
- [50]. Waindi, E. N., Gombe, S. and Oduor–Okelo, D. (1986). Plasma testosterone in *Trypanosoma congolense* infected Togara goats. *Archives of Andrology*, 17:9-17
- [51]. Wellde, B. T., Reardon, M. I, Chumo, D. A., Kovatch, R. M., Wacma, D., Wykoff, D. E, Mwangi, J., Boyce, W. L, and Williams, J. S. (1989). Cerebral trypanosomiasis in naturally-infected cattle in Larnbwe Valley, South Nyanza, Kenya. *Annals of Tropical Medicine and Parasitology*, 83:151-160

Abdullahi, M. A "Reproductive Organ Damage of Domestic Ruminants Caused By Animal African Trypanosomiasis: A Review." *IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS)* 10.11 (2017): 67-71.