

## Comparison of Multipurpose Leguminous Fodder Shrubs in the Western Middle Hills of Nepal

Ram Prasad Ghimire<sup>1</sup>, Rita Amgain<sup>1</sup>, Durga Devkota<sup>2</sup>, Devi Prasad Adhikari<sup>1</sup>,  
Bala R Bhatta<sup>2</sup> and Naba Raj Devkota<sup>3</sup>

<sup>1</sup>Nepal Agricultural Research Council, Nepal

<sup>2</sup>Agriculture and Forestry University, Rampur, Chitwan, Nepal

<sup>3</sup>Gandaki University, Pokhara, Nepal

---

### Abstract

Many multipurpose perennial leguminous fodder shrubs are introduced to mitigate the feed deficit situations of ruminants in Nepal. To evaluate these shrub species, a study was undertaken in the middle hill ecology of Nepal for two years. The experiment was executed in the Randomized Complete Block Design (RCBD) with five replications to compare the fodder yield and nutrient constituents and fuelwood yield of *Flemingia macrophylla* (Willd.) Merrill, *Leucaena leucocephala* (Lam.) de Wit. and *Tephrosia candida* DC. The results of the study showed that the *F. macrophylla* had produced substantially higher ( $p < 0.05$ ) fodder yield in comparison to *L. leucocephala* and *T. candida*, with greater fodder to fuelwood ratio. The nutrient constituents were superior for the *L. leucocephala*. The fuelwood yield was better for the *T. candida* and *F. macrophylla*. The study had revealed that the *F. macrophylla* shrub could be the better alternative for improving the legume fodder productivity in the Nepali middle hill farming systems, and mixing it with *L. leucocephala* could be a better strategy for further improvement of the fodder quality.

**Key words:** Chemical constituents, defoliation, dried fodder yield, fuelwood

---

Date of Submission: 16-10-2021

Date of Acceptance: 31-10-2021

---

### I. Introduction

Hundreds of fodder species and cultivars are introduced in Nepal in order to mitigate the feed and nutrient deficit situations of ruminants in the country. Among them, several annual, biennial and perennial fodder species of different life forms (herbaceous, creepers, shrubs and tree fodder) are established in the Nepali farming systems [1, 2, 3]. The multipurpose shrubs; *F. macrophylla*, *L. leucocephala* and *T. candida*, are some of those introduced perennial fodder legumes which are most preferred by the farmers in the middle hills and Terai regions of Nepal [4, 5, 6]. *Flemingia macrophylla*, recognized as a promising perennial fodder legume and recommended by several studies for Nepali farming systems [4, 7, 8], is being popular in recent years [9]. Likely, *L. leucocephala* is another commonly grown fodder legume for hedgerow plantations in recent years in Nepal. It is another promising perennial fodder legume with the good fodder yield and quality [1, 3]. Similarly, *T. candida* is also a potential perennial leguminous fodder shrub, which is getting popularity in Nepalese farming systems in the past few years [9, 10, 11].

Limited land holdings and resources for intensive fodder cultivation characterize the hill farming systems in Nepal. In this context, promotions of the perennial leguminous fodder shrubs with the selection of the most suitable species is the felt-need of the recent ruminant production systems in those areas [4, 9]. Due to the scanty information about most appropriate species and inadequate region-specific researches on these fodder legumes, the current attempts to promote these species is not being effective [3, 4, 9]. In this context, a study was undertaken for two years in order to select the better perennial leguminous fodder shrub for the western middle hills of Nepal.

### II. Materials and Methods

The experiment was conducted from April 2013 to March 2015 for two years in the 9-months old mature stands of perennial multipurpose leguminous fodder shrubs at the Sarpani experimental site of Goat Research Station situated in the western middle hills of Nepal. Different species of perennial leguminous fodder shrubs were taken as the treatments; *F. macrophylla*, *L. leucocephala* and *T. candida*. The experiment was laid out in Randomized Complete Block Design (RCBD) with five replications. The planting geometry was

maintained at 0.9 m × 0.7 m for all the shrub species. The defoliation height and the defoliation interval were maintained as 1 m and 12 weeks, respectively for all the shrub species.

The collected fodder samples were analyzed at the laboratory of Animal Nutrition Division, Khumaltar, Lalitpur by using methods of AOAC [12] and Goering and Van Soest [13]. Data was analyzed by using the STAR Statistical Package [14]. The combined analysis of both year data was done by using ANOVA and the year effect was partitioned.

$$\gamma_{ijk} = \mu + \tau_i + c_j + \beta_k + \epsilon_{ijk}$$

Where,  $\mu$  is the mean,  $\tau_i$  is  $i^{th}$  treatment effect,  $c_j$  is the temporal effect,  $\beta_k$  is  $j^{th}$  block effect and  $\epsilon_{ijk}$  is random error of observation

### III. Results

#### Dried fodder yield

The temporal effect (year effect) was obtained significant ( $p > 0.05$ ) to the fodder yield of the leguminous shrub species, although, the interactions between year and species, and year and blocks were obtained non-significant ( $p > 0.05$ ).

The results of the study showed that the annual dried fodder yield of *F. macrophylla* was significantly higher ( $p < 0.05$ ) than *L. leucocephala* and *T. candida* (Table 1). It was higher ( $p < 0.05$ ) for the *F. macrophylla* in comparison to *L. leucocephala* and *T. candida* during every defoliation (first, second, third and fourth defoliations), except *et par* ( $p > 0.05$ ) fodder yield with *T. candida* in the case of third defoliation.

**Table 1.** Dried fodder yield of different perennial leguminous fodder shrubs

Shrub species	FD (June 26), t ha <sup>-1</sup>	SD (Sept 20), t ha <sup>-1</sup>	TD (Dec 14), t ha <sup>-1</sup>	FoD (March 8), t ha <sup>-1</sup>	Total fodder yield, t ha <sup>-1</sup> year <sup>-1</sup>
<i>F. macrophylla</i>	5.73±0.92 <sup>a</sup>	4.29±0.79 <sup>a</sup>	2.77±0.53 <sup>a</sup>	2.56±0.43 <sup>a</sup>	15.35±2.37 <sup>a</sup>
<i>L. leucocephala</i>	2.92±0.31 <sup>b</sup>	3.13±0.56 <sup>b</sup>	1.17±0.26 <sup>b</sup>	1.69±0.21 <sup>b</sup>	8.91±1.46 <sup>b</sup>
<i>T. candida</i>	3.42±1.06 <sup>b</sup>	3.11±1.46 <sup>b</sup>	2.26±0.88 <sup>a</sup>	1.92±0.46 <sup>b</sup>	10.7±3.88 <sup>b</sup>

FD= First defoliation, SD= Second defoliation, TD= Third defoliation, FoD= Fourth defoliation

The mean values in a column with different superscripts is significantly different at 5% probability level

In the case of fodder quality, the *L. leucocephala* fodder had significantly higher ( $p < 0.05$ ) crude protein (CP) content than the fodder of *F. macrophylla* and *T. candida* (Figure 1). The neutral detergent fiber (NDF) and acid detergent fiber (ADF) contents were higher ( $p < 0.05$ ) for *F. macrophylla*. The NDF content was lowest for *T. candida*, followed by *L. leucocephala*. Likewise, the ADF content was statistically similar ( $p > 0.05$ ) for the *T. candida* and *L. leucocephala*.




Figure 1. The crude protein and fiber contents in the fodders of different perennial leguminous shrubs in western middle hills of Nepal

Similarly, the calcium content was found significantly higher ( $p < 0.05$ ) for *T. candida* followed by *L. leucocephala* and *F. macrophylla* fodders, although, all the species had good mineral contents (Figure 2). Likely, *T. candida* and *L. leucocephala* fodders had better ( $p < 0.05$ ) phosphorus contents than the *F. macrophylla* in the study.




Figure 2. The calcium and phosphorus contents in the fodders of different perennial leguminous shrubs in western middle hills of Nepal

In the study, *L. leucocephala* and *T. candida* had produced statistically similar ( $p>0.05$ ) total annual fuelwood yield, which were significantly higher ( $p<0.05$ ) than the fuelwood yield of *F. macrophylla* (Table 2). The greater variations in the fuelwood yield among the species were obtained during the first defoliation.

Table 2. Fuelwood yield (on fresh basis) of different perennial leguminous fodder shrub

Shrub species	FD (June 26), t ha <sup>-1</sup>	SD (Sept 20), t ha <sup>-1</sup>	TD (Dec 14), t ha <sup>-1</sup>	FoD (March 8), t ha <sup>-1</sup>	Total fodder yield, t ha <sup>-1</sup> year <sup>-1</sup>
<i>F. macrophylla</i>	4.01±0.47 <sup>b</sup>	4.62±0.34	2.36±0.20 <sup>c</sup>	2.43±1.12 <sup>b</sup>	13.42±3.04 <sup>b</sup>
<i>L. Leucocephala</i>	5.76±2.26 <sup>a</sup>	4.37±1.14	3.28±1.28 <sup>b</sup>	3.46±1.16 <sup>a</sup>	16.87±5.64 <sup>a</sup>
<i>T. candida</i>	5.38±1.87 <sup>a</sup>	4.46±1.89	4.01±1.92 <sup>a</sup>	2.28±0.34 <sup>b</sup>	16.13±6.13 <sup>a</sup>

FD= First defoliation, SD= Second defoliation, TD= Third defoliation, FoD= Fourth defoliation

The mean values with different superscripts in a column is significantly different at 5% probability level

The green fodder to fuelwood ratio was higher for *F. macrophylla* followed by *T. candida* and *L. leucocephala*. The ratio for *F. macrophylla* was gradually decreased to later defoliations. It was increased during September and March defoliations in the case of *T. candida* and *L. leucocephala*.




Figure 3. The green fodder to fuelwood ratio (by fresh weight) in different perennial leguminous fodder shrubs in western middle hills of Nepal

#### IV. Discussion

The significant temporal effect on fodder yield, but non-significant interaction effects of year with fodder yield, and year with the block had indicated the consistent yield variations due to the years over different species and the blocks, although, the fodder yield of the individual shrub was varied between first and second years. The reason might be associated with the year-to-year variations on plant environment such as temperature, moisture, sunshine, soil nutrients and other abiotic factors [15].

The results of the study- *F. macrophylla* had produced consistently higher dried fodder yield with a greater fodder to fuelwood ratio (by fresh weight basis) compared to *T. candida* and *L. leucocephala* in all defoliations in the study- is supported by the report of another study [16]. The authors reported that the *F. macrophylla* had yielded a significantly higher annual weight of leaves and twigs (35 t ha<sup>-1</sup>) compared to *L. leucocephala* (27 t ha<sup>-1</sup>) during the plantain alley cropping. In the study, the data of the dried fodder yield had shown that the variations within the yield of a shrub was higher for *T. candida* compared to the data of *F. macrophylla* and *L. leucocephala*. This larger dispersion of the data indicated higher variability in the fuelwood yield of *T. candida* among different plants, blocks and years.

The CP content obtained for *L. leucocephala* fodder in this study were within the ranges reported by different authors, Topps [17] and Garcia et al., [18], which was significantly higher than *F. macrophylla* and *T. candida*. Nevertheless, the CP values obtained for all the fodder legumes were well above the minimum requirements of 11.9 percent and 12.4 percent recommended for growing and lactating 400 kg cow [19]. It indicates the adequacy of CP levels in all the evaluated fodder species to the ruminant nutrition in Nepali middle hills, although, the *F. macrophylla* had a significantly lower fraction of CP compared to *L. leucocephala* and *T. candida*.

*Flemingia macrophylla* is a perennial leguminous fodder that is characterized by the high fiber content in leaf therefore lower digestibility [8]. In this study too, the fodder of *F. macrophylla* had significantly higher NDF (which is a measure of the plant cell wall material) and ADF contents with the values of 53.89 and 46.78 percentages, respectively that might reduce the fodder quality in many cases. The high ADF value of *F. macrophylla* was probably due to the high cellulose content of the fodder. The finding of this study is in agreement with the reports of other authors who described that the nutritive quality of *F. macrophylla* fodder is lower than that of *L. leucocephala* and other perennial leguminous fodder shrubs, such as *Gliricidia sepium* (Jacq.) Walp. and *Cratylia argentea* (Desv.) O. Kuntze [20, 21]. *Tephrosia candida* had better NDF and ADF contents than other fodder legumes evaluated. On the other hand, higher NDF and ADF contents of *F. macrophylla* could have contributed to higher dried fodder yield and might have decreased the CP and mineral (calcium and phosphorus) contents of the fodder in comparison to other species. It had shown lower fodder quality while comparing these three perennial leguminous fodder legumes. Significantly higher CP percent on *L. leucocephala* than *T. candida*, and similar NDF and ADF contents in *L. leucocephala* and *T. candida* was obtained in another study in Nigeria [22]. The NDF values were found as 39.8 percent and 41.0 percent in *L. leucocephala* and *T. candida*, respectively whereas ADF fractions were 34.30 percent and 36.80 percent for *L. leucocephala* and *T. candida*, respectively [22]. These values were near to the values obtained in this study.

*F. macrophylla* had consistently shown higher green fodder to fuelwood ratio in all defoliations in the study which was in line with the findings of Perera et al. [23]. The authors reported the high leaf:stem ratio in *F. macrophylla*. The results of the study showed that *F. macrophylla* was more consistent in the fodder and fuelwood yield with lower standard deviations compared to the *L. leucocephala* and *T. candida* in all harvests, and in annual yield during the combined analysis of both years' data. But, *F. macrophylla* had a greater reduction in the fodder to fuelwood ratio from the second defoliation and onwards indicating the greater wooding on the fodder branches during September, December and March defoliations. On the other hand, *L. leucocephala* and *T. candida* had better fodder to fuelwood ratios during post-monsoon (September 20) and spring (March 8) defoliations.

#### V. Conclusion

The study concluded that the *F. macrophylla* had better performed than *L. leucocephala* and *T. candida* in the western middle hills of Nepal in terms of fodder yield and fodder to fuelwood ratio, with satisfactory fodder quality. It is a better perennial leguminous fodder shrub than *L. leucocephala* and *T. candida* to mitigate the feed and nutrient deficit situations in western middle hills of Nepal. Generations of the technologies and preparations of the package of practice (PoPs) of *F. macrophylla* and its promotions in future can make substantial contributions to improve the ruminant productivity in the western middle hills of Nepal.

#### Acknowledgements

The authors are highly thankful to the Goat Research Station (Goat), Nepal Agricultural Research Council for the financial support and providing the research field during the experimentation.

## References

- [1]. Upreti, C. R., & Shrestha, B. (2006). *Nutrient Contents of Fodders and Feeds in Nepal*. Kathmandu, Nepal: Animal Nutrition Division, Nepal Agricultural Research Council.
- [2]. Singh, S. B., & Singh, N. (2019). Nepal livestock feed balance and strategies to address the feed deficit. *Journal of Agriculture and Forestry University*, 3, 159-171.
- [3]. Ghimire, R.P., Devkota, N.R., & Tiwari, M.R. (2013). Seasonal productivity of *Flemingia macrophylla* under different defoliation frequencies. *Global Journal of Science Frontier Research: Agriculture and Veterinary*, 13 (14), 12-18.
- [4]. Ghimire, R. P. (2013). Fodder management strategies and their challenges for commercial goat production in Nepal. In T. B. Gurung, B. R. Joshi, U. M. Singh, B. S. Shrestha, K. P. Rijal & D. R. Khanal (Eds.), *The Proceedings of the National Workshop on Research and Development Strategies for Goat Enterprises in Nepal* (pp. 24-41). Kathmandu, Nepal 27-28 September, 2012. Kathmandu: Nepal Agricultural Research Council, Department of Livestock Services, Heifer Project International-Nepal.
- [5]. Tiwari, S. P., & Yadav, J. P. (2014). Consequences of number of irrigations and their interval on seed yield and biomass production of berseem in Nepal. In Muhammad, D., Misri, B., E.-N. M, K. S, & Serkan, A. (Eds.), *Egyptian Clover (Trifolium alexandrinum), King of forage crops* (pp. 95-97). Retrieved from <http://www.fao.org/3/a-i3500e.pdf>.
- [6]. Osti, N. P. (2020). Animal feed resources and their management in Nepal. *ACTA Scientific Agriculture*, 4 (1), 2-14.
- [7]. Singh, S.B. (2000). Strategies and alternative approaches for future growth of livestock sector in Nepal. *Proceedings of the 3<sup>rd</sup> National Animal Science Convention* (pp. 1-14), Aug 27-28, 1997, Kathmandu, Nepal: Nepal Animal Science Association.
- [8]. Ghimire, R. P. (2021). Fodder and pasture research in Nepal: Status, challenges and way forward. In: Shrestha, S.P. (Ed.). *Proceedings of 12<sup>th</sup> National Workshop on Livestock and Fisheries Research in Nepal* (pp. 01-13), 3-4 March 2021. Rampur, Chitwan: Nepal Agricultural Research Council, Kathmandu, Nepal.
- [9]. NAFLQML. (2019). *Balance sheet of animal feed and forage seed of Nepal and impact study of forage mission program*. National Animal Feed and Livestock Quality Management Laboratory, Hariharbhawan, Lalitpur
- [10]. Amatya, S.M., Pandit, B., Subedi, Y.R., Nuberg, I., & Shrestha, K. (2014). Survey of Agroforestry systems in Kabhre and Lamjung districts of Nepal. *Research Paper Series on Agroforestry and Community Forestry in Nepal (2014-06)*, pp. 1-21.
- [11]. Gaire, S.P. (2015). Enhancing livelihood of poor farmers through the promotion of sustainable goat husbandry. *Project Compilation Report* (pp. 103-106). National Agricultural Research and Development Fund, Kathmandu, Nepal.
- [12]. AOAC. (1980). *Official Methods of Analysis of the Association of Official Analytical Chemist* (13th ed.). Washington DC: Association of Official Analytical Chemists.
- [13]. Goering, H. K., & Van Soest, P. J. (1970). Fodder fiber analysis. *USDA Agriculture Handbook* (No. 379.) Washington DC: USDA, ARS.
- [14]. STAR. (2014). *Statistical Tool for Agricultural Research*. Los Banos, Laguna: Biometrics and Breeding Informatics, PBGB Division, International Rice Research Institute.
- [15]. Alam, M.R. (2005). Environmental factors affecting yield and nutrient composition of forages. In: Impact of global change on availability of fodder and forage and performance of livestock in South Asia: *Final Report for APN project 2005-24-NSG-Babar* (pp. 1-18). Asia-Pacific Network for Global Change Research.
- [16]. Banful, B., Dzieror, A., Ofori, I., & Hemeng, O.B. (2000). Yield of plantain alley cropped with *Leucaena leucocephala* and *Flemingia macrophylla* in Kumasi, Ghana. *Agroforestry Systems*, 49, 189-199.
- [17]. Topps, J. H. (1992). Potential, composition and use of legume shrubs and trees as fodder or livestock in the tropics- a review. *Journal of Agricultural Science*, 118, 1-18
- [18]. Garcia, G.W., Ferguson, T.U., Neckles, F.A., & Archibald, K.A.E. (1996). The nutritive value and forage productivity of *Leucaena leucocephala*. *Animal Feed Science Technology*, 60, 29-41.
- [19]. NRC (National Research Council). (1981). *Effect of environment on nutrient requirement of domestic animals*. Washington, DC: National Academy Press. Retrieved from [http://books.nap.edu/openbook.php?record\\_id=4963&page=R1](http://books.nap.edu/openbook.php?record_id=4963&page=R1).
- [20]. Shelton, H.M. (2000). Tropical forage tree legumes in agroforestry systems. *Unasylva*, 51, 25-32.
- [21]. Anderson, M.S. (2006). *Diversity in the tropical multipurpose legumes Cratylia argentea* (Desv.) O. Kuntze and *Flemingia macrophylla* (Willd.) Merrill (Doctoral Thesis), University of Hohenheim, 176p.
- [22]. Odedire, J.A., & Babayemi, O.J. (2007). Preliminary study on *T. candida* as forage alternative to *L. leucocephala* for ruminant nutrition on Southeast Nigeria. *Livestock Research for Rural Development*, 19 (9). Retrieved from <http://www.lrrd.org/lrrd19/9/oded19128.htm>.
- [23]. Perera, A.N.F. (1994). Yield and feeding value of leguminous fodder species grown as hedgerows in slopy lands of Sri Lanka. *Proceedings of the 7<sup>th</sup> AAAP Animal Science Congress*, 11-16 July, 1994. Bali, Indonesia, 2:171-172

Ram Prasad Ghimire, et. al. "Comparison of Multipurpose Leguminous Fodder Shrubs in the Western Middle Hills of Nepal." *IOSR Journal of Research & Method in Education (IOSR-JRME)*, 11(05), (2021): pp. 24-28.